arXiv:math-ph/0501042v1 14 Jan 2005

Finsleroid–Relativistic Time–Asymmetric Space and Quantized Fields

G. S. Asanov

Division of Theoretical Physics, Moscow State University
119992 Moscow, Russia
(e-mail: asanov@newmail.ru)

For well-defined Finsleroid-relativistic space $\mathcal{E}_g^{SR}$ (with the upperscript SR meaning Special-Relativistic) due only to accounting a characteristic parameter g which measures the deviation of the geometry from its pseudoeuclidean precursor, the creation of the respective quantization programs for relativistic physical fields seems to be an urgent task. The parameter may take on the values over all the real range; at $g = 0$ the space is reduced to become an ordinary pseudoeuclidean one. In the present work, the formulation of theory for relativistic physical fields in such a space is initiated. A general method to solve respective scalar, electromagnetic, and spinor field equations is proposed basing on the conformal flatness. At any value of the parameter, the expansion of the relativistic fields with respect to non-plane waves appeared is found, which proposes a base upon which the fields can be quantized in context of the Finsleroid-relativistic approach. Remarkably, the regulators can naturally be proposed to overcome divergences in relativistic field integrals. The respective key and basic concepts involved are presented. For short, the abbreviations FMF, FMT, and FHF will be used for the Finsleroid-relativistic metric function, the associated metric tensor, and the associated Hamiltonian function, respectively.

1. Introduction

Choosing the basic metric function to be of the pseudoeuclidean form is the geometric base proper for the nowadays relativistic field theory. The Finsler-geometry methods [1-2] may suggest the possibility of extending the theory by using metric functions of essentially more general types. Below, we develop the Finslerian $\mathcal{E}_g^{SR}$–space approach specified by a characteristic parameter g. Conceptually, our interpretation of the possibility is based upon treating the centered $\mathcal{E}_g^{SR}$–space as an Observation Space ($\mathcal{OS}$ for short) for the quantum processes. Some value of g is implied to be attached to $\mathcal{OS}$.

The notation R will be used for the four-dimensional vectors that issue from the centre point "O" and belong to the space: $R \in \mathcal{OS}$. The parameter g is independent of the vector argument R (is a constant over the whole $\mathcal{OS}$). For the sake of convenience in process of calculations, the space $\mathcal{OS}$ or $\mathcal{E}_g^{SR}$ will be referred to a rectilinear coordinate system, which centre being identified with the "O", and the components of the vectors with respect to the system will be used: $R = \{R^p\} = \{R^0, R^1, R^2, R^3\} \equiv \{T, X, Y, Z\}$.

We use a particular special-relativistic FMF $F = F(g; T, X, Y, Z)$ subject to the following attractive conditions:

(P1) *The indicatrix-surface* $\mathcal{I}_g$ *defined by the equation* $F(g; T, X, Y, Z) = 1$ *is a regular space of a constant negative curvature,* R_{Ind},

which is a convenient stipulating to seek for the nearest Riemannian-to-Finslerian relativistic generalizations;

(P2) *The FMF is compatible with the principle of spatial isotropy* (*P–parity*);

(P3) *The associated FMT is of the time-space signature* $(+ - - -)$;

(P4) *The principle of correspondence holds true*,

that is, the associated FMT reduces exactly to its ordinary known special– or general–relativistic prototype when $R_{\text{Ind}} \to -1$, which physical significance is quite transparent. The limit value $R_{\text{Ind}} = -1$ corresponds precisely to $g = 0$, so that $F(0; T, X, Y, Z) = \sqrt{T^2 - X^2 - Y^2 - Z^2}$.

All the items (P1)–(P4) are obeyed whenever one makes the choice (A.12) (see Appendix A). Vice versa, we can claim the following

THE UNIQUENESS THEOREM. *The properties* (P1)–(P4), *when treated as conditions imposed on the FMF, specify it unambiguously in the form given by the definition* (A.12).

The indicatrix curvature value is given by (A.29).

NOTE. The $\mathcal{E}_g^{SR}$–space is the Finsleroid-type relativistic space founded upon retaining the P–parity at the expense of violating the T–parity. The resultant FMF F proves to be *T-asymmetric*. The associated FMT exhibits the 6-parametrical invariance. Various aspects of the space were formulated earlier [3-5]. There exists an alternative space in which the T–parity holds under violating the P– parity, assuming existence of a preferred geometrically-distinguished spatial direction [6].

The basic properties of the $\mathcal{E}_g^{SR}$–space approach to the theory of relativistic fields can readily be seen and exercised in case of the scalar field (Section 2). A novel qualitative feature is that the conservation laws cease being integrable, for they involve now the connection coefficients $C_p{}^q{}_r$ (see (2.17)–(2.18)). In Section 3, the electromagnetic field is tackled with. In Section 4, the spinor field equations are generalized accordingly. In analyzing the field equations obtained, it is convenient to convert the consideration into the quasi-pseudoeuclidean space. Solutions to the equations prove to be direct images of solutions to the equations for quasi-pseudoeuclidean fields. In the $O_1(g)$–approximation (which is accounting for the Finslerian corrections proportional to degrees of the characteristic parameter g of the order not higher than the first one) the latter equations are of the ordinary pseudoeuclidean forms, so that for them the ordinary relativistic plane- wave expansion can be applied. The fact is that in the $O_1(g)$–approximations the form of the quasi-pseudoeuclidean metric tensor is merely of the pseudoeuclidean type (corrections are proportional to g^2).

However, the possibility of solving the $\mathcal{E}_g^{SR}$–space relativistic field equations in the general case not adhering at the $O_1(g)$–approximation is not obvious. In Section 5 the method is proposed to rigorously solve the $\mathcal{E}_g^{SR}$–space electromaganetic field equations basing upon the remarkable fact that the respective Lagrangian density is conformally invariant. In Section 6 we show how the Lagrangians for scalar and spinor fields can be extended to entail upon appropriate changes of variables the field equations of ordinary pseudoeuclidean form. To this end, the concepts of conformal scalar and spinor fields are introduced. Therefore, we obtain the total method to solve the $\mathcal{E}_g^{SR}$–equations for all the electromagnetic, scalar, and spinor fields. Particularly, the $\mathcal{E}_g^{SR}$–space versions of the Coulomb law and the Yakawa potential can explicitly be proposed.

Also, we find relevant extensions of plane-wave decompositions (as well as of spherical-type solutions). This circumstance opens up due vistas to quantize $\mathcal{E}_g^{SR}$–space fields in a convenient and rigorous way (Section 7). The waves are not plane but conformally plane and the expansion of fields with respect to such waves is offered. The operator of the four-dimensional momentum can readily be indicated for such waves.

The $\mathcal{E}_g^{SR}$–space approach proposed in the present work may, in principle, provide one with simple explanation of the nature of ultraviolet as well as angular divergences (which are characteristic of ordinary Lorentz-invariant theory of quantized fields), for the limiting transition with $g \to 0$ in the $\mathcal{E}_g^{SR}$–space extended integrals to the integrals of the Lorentz-invariant theory proves to be far from being always possible! There appear convenient regulators, the $\hat{\mathcal{E}}_g^{SR}$–space weights, to perform non-singular integrations over momenta (Section 8).

Appendices A and B are devoted to explaining the basic properties of the $\mathcal{E}_g^{SR}$–space. Appendix C describes and visualizes the behaviour of fronts of respective non-plane waves. Appendices D and E present the conformal properties and various $O_1(g)$–approximations.

The pseudoeuclidean metric tensor will be presented by the components e_{ij}, so that $e_{ij} = e^{ij} = \mathrm{diag}(1, -1, -1, -1)$; the short derivative designations $\partial_i = \partial/\partial t^i$ and $\partial_p = \partial/\partial R^p$ will be used.

2. $\mathcal{E}_g^{SR}$–space extension of Klein-Gordon equation

Let $\phi = \phi(R)$ be a complex scalar field, so that

$$\phi(R) = \phi_1(R) + i\phi_2(R), \qquad \phi^*(R) = \phi_1(R) - i\phi_2(R), \tag{2.1}$$

where $\phi_1(R)$ and $\phi_2(R)$ are two real scalar functions; the star (*) means the complex conjugation. Introducing also the notation $\phi_p = \partial_p \phi$ and $\phi_p^* = \partial_p \phi^*$ for partial derivatives with respect to the argument R^p, we construct the Lagrangian

$$L_\phi = g^{pq}(g;R)\phi_p^*(R)\phi_q(R) - m^2\phi^*(R)\phi(R) \tag{2.2}$$

(m denotes the rest mass of the scalar particle) and the Lagrangian density

$$\Lambda_\phi = L_\phi J, \tag{2.3}$$

where

$$J \ := \sqrt{|\det(g_{pq}(g;R))|} = (j(g;R))^4 \tag{2.4}$$

denotes the Jacobian for the FMT (see (A.18)), to use the action integral

$$S\{\phi\} = \int \Lambda_\phi d^4R. \tag{2.5}$$

The associated Euler-Lagrange derivatives

$$\mathcal{E}_\phi \ := \partial_q\frac{\partial\Lambda_\phi}{\partial\phi_q} - \frac{\partial\Lambda_\phi}{\partial\phi}, \qquad \mathcal{E}_\phi^* \ := \partial_q\frac{\partial\Lambda_\phi}{\partial\phi_q^*} - \frac{\partial\Lambda_\phi}{\partial\phi^*} \tag{2.6}$$

can be found from Eqs. (2.1)–(2.6) to be

$$\frac{1}{J}\mathcal{E}_\phi = \Box\phi^* + m^2\phi^*, \qquad \frac{1}{J}\mathcal{E}_\phi^* = \Box\phi + m^2\phi, \tag{2.7}$$

where

$$\Box = \frac{1}{J}\partial_p\Big(Jg^{pq}\partial_q\Big). \tag{2.8}$$

Since $\delta S = 0 \implies \mathcal{E}_\phi = 0$, we get the following $\mathcal{E}_g^{SR}$*–space scalar field equation*:

$$\Box\phi + m^2\phi = 0. \tag{2.9}$$

The respective $\mathcal{E}_g^{SR}$*–space scalar field current density*

$$\mathcal{J}_p \ := i(\phi^*\partial_p\phi - \phi\partial_p\phi^*)J \tag{2.10}$$

is conserved over solutions to the equation (2.9), according to

$$\partial_p\mathcal{J}^p = 0, \tag{2.11}$$

where $\mathcal{J}^p = g^{pq}\mathcal{J}_q$. Therefore the charge is still conserved in the ordinary sense:

$$Q = \int \mathcal{J}^0 d^3\mathbf{R} \tag{2.12}$$

and

$$\partial_0 Q = -\int \mathcal{J}^a d\Sigma_a. \tag{2.13}$$

Considering the respective energy-momentum tensor of scalar field:

$$T_p{}^q \ := (\phi_p\frac{\partial\Lambda_\phi}{\partial\phi_q} + \phi_p^*\frac{\partial\Lambda_\phi}{\partial\phi_q^*} - \delta_p^q\Lambda_\phi)/J, \tag{2.14}$$

we obtain

$$T_p{}^q = \phi_p \phi^{*q} + \phi^*_p \phi^q - \delta^q_p L_\phi, \tag{2.15}$$

where

$$\phi^q(R) = g^{qp}(g;R)\phi_p(R), \qquad \phi^{*q}(R) = g^{qp}(g;R)\phi^*_p(R). \tag{2.16}$$

Now, the $\mathcal{E}_g^{SR}$–space extension of the conservation law for the scalar field energy-momentum tensor reads

$$\mathcal{D}_q T_p{}^q \ := \partial_q T_p{}^q - C_p{}^r{}_q T_r{}^q + C_t{}^q{}_q T_p{}^t \tag{2.17}$$

and

$$\mathcal{D}_q T_p{}^q = 0, \tag{2.18}$$

what can be verified by direct calculations; $C_p{}^r{}_q$ are coefficients (A.20).

Let us apply the quasi-pseudoeuclidean transformation $t^i = \sigma^i(g;R)$ (see (B.1)–(B.2)) and introduce the transformed scalar field in accordance with $\phi(R) = u\left(\sigma(g;R)\right)$, so that

$$u(t) = \phi(R). \tag{2.19}$$

We call the field $u(t)$ the *quasi-pseudoeuclidean image of the scalar field* $\phi(R)$. We get

$$\Lambda_\phi = \Lambda_u, \tag{2.20}$$

$$L_u = n^{ij}(g;t)u^*_i(t)u_j(t) - m^2 u^*(t)u(t), \tag{2.21}$$

and

$$\Lambda_u = L_u h^{-3}, \tag{2.22}$$

where the equality (B.19) has been used; the tensor $n^{ij}(g;t)$ is given by (B.16). The action integral (2.5) goes over in

$$S\{u\} = \int \Lambda_u d^4 t. \tag{2.23}$$

From $\delta S = 0 \implies \mathcal{E}_u = 0$ we obtain the *quasi-pseudoeuclidean scalar field equation*:

$$\Box u + m^2 u = 0, \tag{2.24}$$

where

$$\Box = \partial_i (n^{ij}\partial_j). \tag{2.25}$$

Considering the *quasi-pseudoeuclidean image of the density of four-dimensional Finslerian scalar current*

$$\mathcal{J}_i \ := i(u^*\partial_i u - u\partial_i u^*)h, \tag{2.26}$$

we arrive at the conclusion that the current is conserved over solutions to the equations (2.24):

$$\partial_i \mathcal{J}^i = 0, \tag{2.27}$$

where $\mathcal{J}^i = n^{ij}\mathcal{J}_j$.

The energy-momentum tensor

$$T_i{}^j \ := \Big[u_i \frac{\partial \Lambda_u}{\partial u_j} + u^*_i \frac{\partial \Lambda_u}{\partial u^*_j} - \delta^j_i \Lambda_u\Big]/h \tag{2.28}$$

can be rewritten as

$$T_i^j = u_i(t)u^{*j}(t) + u_i^*(t)u^j(t) - \delta_i^j L_u, \tag{2.29}$$

where

$$u^j(t) = n^{ji}(g;t)u_i(t), \qquad u^{*j}(t) = n^{ji}(g;t)u_i^*(t). \tag{2.30}$$

From (2.24) the following conservation law follows for the quasi-pseudoeuclidean tensor (2.29):

$$\mathcal{D}_j T_i^j \ := \partial_j T_i^j - N_i{}^m{}_j T_m^j + N_n{}^j{}_j T_i^n \tag{2.31}$$

and

$$\mathcal{D}_j T_i^j = 0, \tag{2.32}$$

where $N_i{}^m{}_j$ — the connection coefficients of the quasi-pseudoeuclidean space (see (B.26)).

Thus we have proved

PROPOSITION 1. *Solutions to the Finslerian scalar field equations* (2.9) *are the quasi-pseudoeuclidean images of solutions to the equations* (2.24) *for quasi-pseudoeuclidean scalar field.*

The transformations

$$\mathcal{J}^p(R) = R_i^p(g;t)\mathcal{J}^i(t) \tag{2.33}$$

and

$$\mathcal{J}^i(t) = t_p^i(g;R)\mathcal{J}^p(R) \tag{2.34}$$

together with

$$T^{pq}(R) = R_i^p(g;t)R_j^q(g;t)T^{ij}(t) \tag{2.35}$$

and

$$T^{ij}(t) = t_p^i(g;R)t_q^j(g;R)T^{pq}(R) \tag{2.36}$$

are implied; t_p^i and R_i^p are the coefficients (B.9) and (B.13).

In the $O_1(g)$*–approximation* we have simply $n_{ij} = e_{ij}$ and $n^{ij} = e^{ij}$. The coefficients $N_i{}^k{}_j$ are proportional to g^2 (see (B.26)) and, therefore, disappear at the $O_1(g)$–level. With these restrictions, the equation (2.24)–(2.25) reduces to the ordinary Klein-Gordon form, so that we are entitled to apply the ordinary plane-wave expansion to the quasi-pseudoeuclidean image of scalar field:

$$u(t) = \frac{1}{(2\pi)^{3/2}} \int e^{ik_j t^j} u(k)\delta(k^2 - m^2)d^4k \tag{2.37}$$

and

$$u(t) = u^+(t) + u^-(t) \tag{2.38}$$

with

$$u^+(t) = \frac{1}{(2\pi)^{3/2}} \int e^{ik_j t^j} u^+(k)\delta(k^2 - m^2)d^4k, \qquad k_0 > 0, \tag{2.39}$$

$$u^-(t) = \frac{1}{(2\pi)^{3/2}} \int e^{-ik_j t^j} u^-(k)\delta(k^2 - m^2)d^4k, \qquad k_0 > 0, \tag{2.40}$$

$$(u^+(k))^* = u^{*-}(k), \qquad (u^-(k))^* = u^{*+}(k), \tag{2.41}$$

$$u^+(\mathbf{k}) = \frac{u^+(k)}{\sqrt{2k_0}}, \qquad u^-(\mathbf{k}) = \frac{u^-(k)}{\sqrt{2k_0}}, \tag{2.42}$$

$$u^+(t) = \frac{1}{(2\pi)^{3/2}} \int e^{ik_j t^j} u^+(\mathbf{k}) \frac{d\mathbf{k}}{\sqrt{2k_0}}, \tag{2.43}$$

$$u^-(t) = \frac{1}{(2\pi)^{3/2}} \int e^{-ik_j t^j} u^-(\mathbf{k}) \frac{d\mathbf{k}}{\sqrt{2k_0}}, \tag{2.44}$$

where $k_0 = \sqrt{\mathbf{k}^2 + m^2} > 0$. The energy-momentum conservation law (2.31)–(2.32) takes on the ordinary integrable form

$$\partial_j T_i^j = 0. \tag{2.45}$$

From the above we obtain

PROPOSITION 2. *At the $O_1(g)$–approximation level of consideration the quasi-pseudoeuclidean image* (2.19) *of scalar field is reduced to merely pseudoeuclidean case.*

Returning back from the quasi-pseudoeuclidean space to the initial $\mathcal{E}_g^{SR}$–space, we observe that the plane-wave decomposition should be replaced by the non-plane-wave decomposition:

$$\phi(R) = \phi^+(R) + \phi^-(R), \qquad \phi^*(R) = \phi^{*+}(R) + \phi^{*-}(R) \tag{2.46}$$

with the functions

$$\phi^+(R) = \frac{1}{(2\pi)^{3/2}} \int e^{ik_n \sigma^n(g;R)} \phi^+(k) \delta\left(k^2 - m^2\right) d^4k \tag{2.47}$$

and

$$\phi^-(R) = \frac{1}{(2\pi)^{3/2}} \int e^{-ik_n \sigma^n(g;R)} \phi^-(k) \delta\left(k^2 - m^2\right) d^4k. \tag{2.48}$$

The ordinary conjugation properties remain valid:

$$(\phi^+(k))^* = \phi^{*-}(k), \qquad (\phi^-(k))^* = \phi^{*+}(k). \tag{2.49}$$

Thus, the $\mathcal{E}_g^{SR}$–space waves of scalar particles are not plane unless $g = 0$. On introducing the functions

$$u^+(\mathbf{k}) = \phi^+(\mathbf{k}) = \frac{\phi^+(k)}{\sqrt{2k_0}}, \qquad u^-(\mathbf{k}) = \phi^-(\mathbf{k}) = \frac{\phi^-(k)}{\sqrt{2k_0}}, \tag{2.50}$$

we get the representations

$$\phi^+(R) = \frac{1}{(2\pi)^{3/2}} \int e^{ik_n \sigma^n(g;R)} \phi^+(\mathbf{k}) \frac{d^3\mathbf{k}}{\sqrt{2k_0}} \tag{2.51}$$

and

$$\phi^-(R) = \frac{1}{(2\pi)^{3/2}} \int e^{-ik_n \sigma^n(g;R)} \phi^-(\mathbf{k}) \frac{d^3\mathbf{k}}{\sqrt{2k_0}}. \tag{2.52}$$

3. $\mathcal{E}_g^{SR}$–space extension of electromagnetic field equations

Let a covariant field $\mathcal{A} = \{A_p(R)\}$ be an electromagnetic vector potential. We shall use the partial derivatives $A_{p,q} = \partial_q A_p$ and, as usually, construct the *tension tensor of electromagnetic field*

$$F_{pq} = A_{p,q} - A_{q,p} \tag{3.1}$$

and introduce the concomitant tensors

$$F_p{}^q = g^{qr}F_{pr}, \qquad F^{pq} = g^{pr}g^{qs}F_{rs}. \tag{3.2}$$

After that, we can introduce the *density of the $\mathcal{E}_g^{SR}$–space induction tensor*

$$\mathcal{D}^{pq}(R) \ := J(g;R)g^{pt}(g;R)g^{qs}(g;R)F_{ts}(R) \tag{3.3}$$

and use the direct extension

$$\Lambda_A = -\frac{1}{4}\mathcal{D}^{pq}F_{pq} \tag{3.4}$$

of the classical (rescaled) Lagrangian density of electromagnetic field to set forth the action integral

$$S\{A\} = \int \Lambda_A d^4R. \tag{3.5}$$

The associated Euler-Lagrange derivatives

$$\mathcal{E}_A^p \ := \partial_q \frac{\partial \Lambda_A}{\partial A_{p,q}} = 2\partial_q \frac{\partial \Lambda_A}{\partial F_{qp}} \tag{3.6}$$

can be found from (3.1)–(3.5) to read

$$\mathcal{E}_A^p = \partial_q \mathcal{D}^{pq}. \tag{3.7}$$

Since $\delta S = 0 \implies \mathcal{E}_A^p = 0$, we obtain the *$\mathcal{E}_g^{SR}$–space electromagnetic field equations*

$$\partial_q \mathcal{D}^{pq} = 0 \tag{3.8}$$

(in vacuum). Additionally the cyclic equations

$$\partial_r F_{pq} + \partial_q F_{rp} + \partial_p F_{qr} = 0 \tag{3.9}$$

are fulfilled. The corresponding *energy-momentum tensor of the $\mathcal{E}_g^{SR}$–space electromagnetic field*

$$T_p{}^q \ := -F_{pr}F^{qr} + \frac{1}{4}\delta_p{}^q F_{st}F^{st} \tag{3.10}$$

will satisfy the conservation law of the form (2.17)–(2.18).

Let us introduce the *quasi-pseudoeuclidean image for the initial electromagnetic vector potential $\mathcal{A}$*:

$$U_i(t) \ := R_i^p(g;t)A_p(\mu(g;t)) \tag{3.11}$$

and, after that, introduce the corresponding definitions for the tension tensor

$$f_{ij}(t) \ := R_i^p(g;t)R_j^q(g;t)F_{pq}(R) \tag{3.12}$$

together with the contravariant tensor

$$f^{ij}(t) \ := t_p^i(g;R)t_q^j(g;R)F^{pq}(R). \tag{3.13}$$

The inverse relations take the form

$$A_p(R) = t_p^i(g;R)U_i(t), \tag{3.14}$$

$$F_{pq}(R) = t_p^i(g;t)t_q^j(g;t)f_{ij}(t), \tag{3.15}$$

$$F^{pq}(R) = R_i^p(g;t)R_j^q(g;t)f^{ij}(t), \tag{3.16}$$

$$f^{ij}(t) = n^{im}n^{jn}f_{mn}(t), \tag{3.17}$$

and the representation

$$f_{mn}(t) = \frac{\partial U_n(t)}{\partial t^m} - \frac{\partial U_m(t)}{\partial t^n} \tag{3.18}$$

is valid.

Next, we introduce the Lagrangian density

$$\Lambda_A = \Lambda_U, \tag{3.19}$$

where

$$\Lambda_U = L_U h^3 \tag{3.20}$$

and

$$L_U = -\frac{1}{4}n^{ij}n^{mn}f_{im}(t)f_{jn}(t) \equiv -\frac{1}{4}f^{im}(t)f_{im}(t), \tag{3.21}$$

so that the action integral (3.5) transforms to

$$S\{U\} = \int \Lambda_U d^4t. \tag{3.22}$$

From the action principle $\delta S = 0$ the equations of the ordinary form

$$\frac{\partial f^{in}}{\partial t^i} = 0 \tag{3.23}$$

ensue. Besides, the counterpart of (3.9) is valid:

$$\frac{\partial f_{mn}}{\partial t^i} + \frac{\partial f_{im}}{\partial t^n} + \frac{\partial f_{ni}}{\partial t^m} = 0 \tag{3.24}$$

The conservation law of the form (2.31)–(2.32) is applicable to the energy-momentum tensor

$$T_i^j = -f^{jm}f_{im} + \frac{1}{4}\delta_j^i f^{mn}f_{mn}. \tag{3.25}$$

Inserting any solution $U_i = U_i(t)$ to the equation (3.23)–(3.24) in the right-hand part of the relation (3.14) yields a field $A_p(R)$, which gives a solution to the initial $\mathcal{E}_g^{SR}$–space electromagnetic field equations (3.8).

In the $O_1(g)$*–approximation* the equations (3.23) reduce to the ordinary Maxwell-type equations:

$$\partial_j f^{ij} = 0 \quad \text{with the functions} \quad f^{ij} = e^{im}e^{jn}f_{mn}, \tag{3.26}$$

and we can use the known solutions formed by plane waves:

$$U_i(t)=\frac{1}{(2\pi)^{3/2}}\int e^{ik_jt^j}U_i(k)\delta(k^2)d^4k, \tag{3.27}$$

or in the form

$$U_i(t)=U_i^+(t)+U_i^-(t) \tag{3.28}$$

with the functions

$$U_i^{\pm}(t)=\frac{1}{(2\pi)^{3/2}}\int e^{\pm ik_jt^j}U_i^{\pm}(\mathbf{k})\frac{d\mathbf{k}}{\sqrt{2k_0}}, \tag{3.29}$$

where $k_0=\sqrt{\mathbf{k}^2}$. Returning into the initial space, we get

$$A_p(R)=A_p^+(R)+A_p^-(R) \tag{3.30}$$

with the functions

$$A_p^{\pm}(R)=t_p^i(g;R)U_i^{\pm}(t)=\frac{1}{(2\pi)^{3/2}}t_p^i(g;R)\int e^{\pm ik_n\sigma^n(g;R)}U_i^{\pm}(\mathbf{k})\frac{d\mathbf{k}}{\sqrt{2k_0}}\,. \tag{3.31}$$

Ordinary plane-wave decomposition (3.27) turn over into non-plane-wave decomposition

$$A_p(R)=\frac{1}{(2\pi)^{3/2}}\int e^{ik_n\sigma^n(g;R)}A_p(R,k)\delta(k^2)d^4k, \tag{3.32}$$

where

$$A_p(R,k)=t_p^i(g;R)U_i(k). \tag{3.33}$$

Whence we have arrived at the conclusion:

PROPOSITION 3. *In the* $O_1(g)$*–approximation, solutions to the initial* $\mathcal{E}_g^{SR}$*–space electcromagnetic field equations* (3.8) *present straightforward images of solutions to the linear equations* (3.26) *of the ordinary Maxwell form.*

4. $\mathcal{E}_g^{SR}$–space extension of spinor field equations

For the four-component spinor $\psi(R)$ of the Dirac type we introduce, in agreement with known methods of general-relativistic approaches, the $\mathcal{E}_g^{SR}$*–space spinor derivatives*

$$D_p\psi(R)=\partial_p\psi(R)-Z_p(g;R)\psi(R), \qquad D_p\bar{\psi}(R)=\partial_p\bar{\psi}(R)+\bar{\psi}(R)Z_p(g;R) \tag{4.1}$$

and, respectively, the $\mathcal{E}_g^{SR}$*–space spinor connection coefficients*

$$Z_p(g;R)=-\frac{1}{8}R^{PQ}{}_p(g;R)(\gamma_P\gamma_Q-\gamma_Q\gamma_P), \tag{4.2}$$

where $R^{PQ}{}_p(g;R)$ — the associated Ricci rotation coefficients (see (A.37)–(A.39)) and the identity

$$D_q\gamma^p=\mathcal{D}_q\gamma^p-Z_q\gamma^p+\gamma^pZ_q=0 \tag{4.3}$$

is fulfilled. The notation γ_P is used for the ordinary Dirac's matrices, so that

$$\gamma_P\gamma_Q+\gamma_Q\gamma_P=2a_{PQ}, \tag{4.4}$$

where $\{a_{PQ}\} = \mathrm{diag}(1,-1,-1,-1)$; we have also introduced the notation

$$\gamma^p(g;R) = g^{pq}(g;R)e^P_q(g;R)\gamma_P, \tag{4.5}$$

where $e^P_q(g;R)$ is the orthonormal invariant frame (see (A.35)). The conjugation operation reads

$$\bar{\psi}(R) = (\psi(R))^+\gamma^0(g;R). \tag{4.6}$$

Therefore, the $\mathcal{E}^{SR}_g$–extended Dirac-type equation reads

$$-i\gamma^p(g;R)D_p\psi(R) + m\psi(R) = 0 \tag{4.7}$$

and the conjugated version reads

$$i(D_p\bar{\psi}(R))\gamma^p(g;R) + m\bar{\psi}(R) = 0; \tag{4.8}$$

here, m stands for the rest mass of the spinor particle. For the corresponding current density

$$\mathcal{J}^p = J\bar{\psi}\gamma^p\psi \tag{4.9}$$

the direct calculations yield

$$\partial_p\mathcal{J}^p = J\left(\partial_p(\bar{\psi})\gamma^p\psi + \bar{\psi}\gamma^p\partial_p\psi + \bar{\psi}(\mathcal{D}_p\gamma^p)\psi\right) = J\left((D_p\bar{\psi})\gamma^p\psi + \bar{\psi}\gamma^p(D_p\psi)\right),$$

where the property (4.3) has been used. The spinor energy-momentum tensor is given by the formula

$$T^{pq} = \frac{i}{2}\left(\bar{\psi}\gamma^pD^q\psi - (D^q\bar{\psi})\gamma^p\psi\right). \tag{4.10}$$

By virtue of the equations (4.1)–(4.8) we can readily infer the conservation laws of the form (2.11)–(2.13) and (2.17)–(2.18).

For the spinor

$$v(t) = \psi(\mu(g;t)), \tag{4.11}$$

which arises on performing a due transformation into quasi-pseudoeuclidean space, we get the equations which are similar to those shown above, namely,

$$D_iv(t) = \partial_iv(t) - T_i(g;t)v(t), \qquad D_i\bar{v}(t) = \partial_i\bar{v}(t) + \bar{v}(t)T_i(g;t) \tag{4.12}$$

and we also get respectively the $\mathcal{E}^{SR}_g$*–space spinor connection coefficients*

$$T_i(g;t) = -\frac{1}{8}R^{PQ}{}_i(g;t)(\gamma_P\gamma_Q - \gamma_Q\gamma_P) \tag{4.13}$$

($R^{PQ}{}_i(g;t)$ stand for the quasi-pseudoeuclidean Ricci rotation coefficients; see (B.34)), from the definition of which the following counterpart of the identity (4.3) ensues:

$$D_j\gamma^i = \mathcal{D}_j\gamma^i - T_j\gamma^i + \gamma^iT_j = 0, \tag{4.14}$$

where

$$\gamma^i(g;t) = n^{ij}(g;t)f^P_j(g;t)\gamma_P \tag{4.15}$$

with $f^P_j(g;t)$ being the frame (B.28)). Also,

$$\bar{v}(t) = (v(t))^+\gamma^0(g;t). \tag{4.16}$$

The Dirac-type equation in the quasi-pseudoeuclidean space takes on the form

$$-i\gamma^i(g;t)D_iv(t)+mv(t)=0, \tag{4.17}$$

and for its conjugation we should write

$$i(D_i\bar{v}(t))\gamma^i(g;t)+m\bar{v}(t)=0. \tag{4.18}$$

For the respective spinor current density

$$\mathcal{J}^i=J\bar{v}\gamma^i v \tag{4.19}$$

and for the respective spinor energy-momentum tensor

$$T^{ij}=\frac{i}{2}(\bar{\psi}\gamma^i D^j\psi-(D^j\bar{\psi})\gamma^i\psi) \tag{4.20}$$

the conservation laws of the form (2.27) and (2.31)–(2.32) hold.

The fact that in the quasi-pseudoeuclidean space the $\mathcal{E}_g^{SR}$–space corrections are of the order of g^2 opens up the direct possibility to solve the considered spinor equations *at the $O_1(g)$–approximation* by using results of ordinary relativistic theory. In particular, we can take solutions to be of the plane-wave type:

$$v(t)=\frac{1}{(2\pi)^{3/2}}\int e^{ik_nt^n}\delta(k^2-m^2)v(k)d^4k, \tag{4.21}$$

or

$$v(t)=v^+(t)+v^-(t) \tag{4.22}$$

with the functions

$$v^{\pm}(t)=\frac{1}{(2\pi)^{3/2}}\int e^{\pm ik_jt^j}v^{\pm}(\mathbf{k})d\mathbf{k}, \tag{4.23}$$

where

$$v^{\pm}(\mathbf{k})=\frac{\theta(k_0)v^{\pm}(k)}{2k^0}. \tag{4.24}$$

On returning back in the initial $\mathcal{E}_g^{SR}$–space, we obtain the representation

$$\psi(R)=\frac{1}{(2\pi)^{3/2}}\int e^{ik_n\sigma^n(g;R)}\delta(k^2-m^2)\psi(k)d^4k. \tag{4.25}$$

5. Conformal method of solution for $\mathcal{E}_g^{SR}$–space electromagnetic field equations

Given a vector field $U_i(t)$ in the quasi-pseudoeuclidean space, we perform the transformation to new components $B_j(r)$ according to the vector law

$$U_i(t)\ :=k_i^jB_j(r) \tag{5.1}$$

which owing to (B.35)–(B.38) reads explicitly as

$$U_i=\frac{1}{h}\xi(B_i+\gamma B_mn_in^m). \tag{5.2}$$

Next, we introduce the tensors

$$B_{ij}\ :=\frac{\partial B_j}{\partial r^i}-\frac{\partial B_i}{\partial r^j}, \tag{5.3}$$

$$B_i{}^j := B_{im} e^{jm}, \qquad B^{ij} := B_n{}^j e^{ni}, \tag{5.4}$$

and apply the transformation law

$$f_{ij} = B_{mn} k_i^m k_j^n, \tag{5.5}$$

where f_{ij} are constructed according to the rule (3.18). The concomitant tensors are $f_i{}^j := f_{im} n^{jm}$ and $f^{ij} := f_n{}^j n^{ni}$.

Due insertions yield explicitly

$$f_{ij} = \frac{1}{h^2}\xi^2\Big[B_{ij} + \gamma h^2 (B_{im} n_j - B_{jm} n_i) n^m\Big], \tag{5.6}$$

$$f_i{}^j = \xi^2\Big[B_i{}^j - \gamma h^2 B^j{}_m n_i n^m + \gamma(2h^2 + \frac{1}{h^2})\Big] B_{jm} n^j n^m, \tag{5.7}$$

and

$$f^{ij} = \xi^2\Big[h^2 B^{ij} - \gamma(\gamma^3 + h^2 + h) n^i B^j{}_m n^m + \gamma(2h^4 + 1) B^i{}_m n^j n^m\Big]. \tag{5.8}$$

The equations

$$\frac{\partial B_{im}}{\partial r^j} + \frac{\partial B_{ji}}{\partial r^m} + \frac{\partial B_{mj}}{\partial r^i} = 0 \tag{5.9}$$

are obviously valid. They entail the electromagnetic field equations (3.24). Also, straightforward calculations show that the equations

$$\frac{\partial B^{ij}}{\partial r^j} = 0 \tag{5.10}$$

would entail the electromagnetic field equations (3.23) for the tensor (5.5).

Thus we have arrived at the following

PROPOSITION 4. The solutions to the electromagnetic field equations in the space $\mathcal{E}_g^{SR}$ are the transforms of the solutions to the ordinary Maxwell equations in accordance with the formulae (3.14)–(3.15) and (5.6)–(5.10).

This result offers a handy way to export the electromagnetic field solutions from the ordinary pseudoeuclidean space to the $\mathcal{E}_g^{SR}$–space under study.

In particular, the ordinary plane-wave solution is now extending as follows:

$$B_m = b_m \, \mathrm{e}^{i\Phi}, \qquad b_m = \text{constants}, \tag{5.11}$$

and

$$U_j = a_j \, \mathrm{e}^{i\Phi}, \qquad a_j = \text{constants}, \tag{5.12}$$

with Φ being the phase (B.51).

Taking for definiteness

$$k_n = (k_0, k_1 = -k_0, 0, 0) \tag{5.13}$$

and

$$b_i = (0,0,0,b_3), \quad b^i = (0,0,0,b^3), \quad b^3 = -b_3, \tag{5.14}$$

we get for the vector potential $A_{\{k\}p}$ the following explicit components:

$$A_{\{k\}0} = \left(\frac{g}{2}(2h-1) - (h-1)k\right) \frac{R^3}{|\mathbf{R}|} \frac{j\varkappa b_3}{L}\, \mathrm{e}^{i\Phi}, \tag{5.15}$$

$$A_{\{k\}1} = \left(h - 1 - \frac{1}{2}gk\right) \frac{R^1R^3}{|\mathbf{R}|^2} \frac{j\varkappa b_3}{L}\, \mathrm{e}^{i\Phi}, \tag{5.16}$$

$$A_{\{k\}2} = \left(h - 1 - \frac{1}{2}gk\right) \frac{R^2R^3}{|\mathbf{R}|^2} \frac{j\varkappa b_3}{L}\, \mathrm{e}^{i\Phi}, \tag{5.17}$$

$$A_{\{k\}3} = \left[L + \left(h - 1 - \frac{1}{2}gk\right) \frac{(R^3)^2}{|\mathbf{R}|^2}\right] \frac{j\varkappa b_3}{L}\, \mathrm{e}^{i\Phi}, \tag{5.18}$$

where

$$L = 1 + gk - k^2 \tag{5.19}$$

and (3.14) has been applied. Contracting yields the simple result

$$R^pA_{\{k\}p} = hj\varkappa R^3 b_3\, \mathrm{e}^{i\Phi}\,. \tag{5.20}$$

The notation

$$k = \frac{R^0}{|\mathbf{R}|} \tag{5.21}$$

and the conformal multipliers $\varkappa$ and ξ (see (B.36) and (B.43)) have conveniently been used.

When

$$R^0 = 0, \tag{5.22}$$

we have

$$k = 0, \quad L = 1, \tag{5.23}$$

$$F\big|_{R^0=0} = (-g_-)^{G_+/2}(g_+)^{-G_-/2}|\mathbf{R}|, \qquad j\big|_{R^0=0} = \left(\frac{-g_-}{g_+}\right)^{-G/4}, \tag{5.24}$$

and the quantity Φ reduces to

$$\Phi_0 = k_0C_2|\mathbf{R}|^{\frac{1}{2}(h-1)}(-\frac{G}{2}|\mathbf{R}| - R^1). \tag{5.25}$$

We obtain

$$A_{\{k\}0}\big|_{R^0=0} = \frac{g}{2}(2h-1)\frac{R^3}{|\mathbf{R}|}|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\, \mathrm{e}^{i\Phi_0}, \tag{5.26}$$

$$A_{\{k\}1}\big|_{R^0=0} = (h-1)\frac{R^1R^3}{|\mathbf{R}|^2}|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\, \mathrm{e}^{i\Phi_0}, \tag{5.27}$$

$$A_{\{k\}2}\big|_{R^0=0} = (h-1)\frac{R^2R^3}{|\mathbf{R}|^2}|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\, \mathrm{e}^{i\Phi_0}, \tag{5.28}$$

$$A_{\{k\}3}\big|_{R^0=0}= \left[1+(h-1)\frac{(R^3)^2}{|\mathbf{R}|^2}\right]|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}\,. \tag{5.29}$$

C and C_2 are constants.

For the contravariant components $A^p_{\{k\}}(R) = g^{pq}(g;R)A_{\{k\}q}(R)$ calculations yield the results

$$A^0_{\{k\}} = (kh-f)\frac{R^3}{|\mathbf{R}|}\frac{b^3}{j\varkappa L}\,\mathrm{e}^{i\Phi}, \tag{5.30}$$

$$A^1_{\{k\}} = (h-(k-g)f-L)\,\frac{R^1R^3}{|\mathbf{R}|^2}\frac{b^3}{j\varkappa L}\,\mathrm{e}^{i\Phi}, \tag{5.31}$$

$$A^2_{\{k\}} = (h-(k-g)f-L)\,\frac{R^2R^3}{|\mathbf{R}|^2}\frac{b^3}{j\varkappa L}\,\mathrm{e}^{i\Phi}, \tag{5.32}$$

$$A^3_{\{k\}} = \left[L+(h-(k-g)f-L)\,\frac{(R^3)^2}{|\mathbf{R}|^2}\right]\frac{b^3}{j\varkappa L}\,\mathrm{e}^{i\Phi}, \tag{5.33}$$

where

$$f = k-\frac{g}{2}. \tag{5.34}$$

When

$$R^0 = 0, \tag{5.35}$$

we obtain

$$A^0_{\{k\}}\big|_{R^0=0}= \frac{g}{2}\frac{R^3}{|\mathbf{R}|}\frac{1}{|\mathbf{R}|^{\frac{1}{2}(h-1)}}Cb^3\,\mathrm{e}^{i\Phi_0}, \tag{5.36}$$

$$A^1_{\{k\}}\big|_{R^0=0}= \left(h-1-\frac{g^2}{2}\right)\frac{R^1R^3}{|\mathbf{R}|^2}\frac{1}{|\mathbf{R}|^{\frac{1}{2}(h-1)}}Cb^3\,\mathrm{e}^{i\Phi_0}, \tag{5.37}$$

$$A^2_{\{k\}}\big|_{R^0=0}= \left(h-1-\frac{g^2}{2}\right)\frac{R^2R^3}{|\mathbf{R}|^2}\frac{1}{|\mathbf{R}|^{\frac{1}{2}(h-1)}}Cb^3\,\mathrm{e}^{i\Phi_0}, \tag{5.38}$$

$$A^3_{\{k\}}\big|_{R^0=0}= \left[1+\left(h-1-\frac{g^2}{2}\right)\frac{(R^3)^2}{|\mathbf{R}|^2}\right]\frac{1}{|\mathbf{R}|^{\frac{1}{2}(h-1)}}Cb^3\,\mathrm{e}^{i\Phi_0}\,. \tag{5.39}$$

The associated electromagnetic field tension tensor can also be found explicitly, namely

$$f_{ij} = i(k_ib_j-k_jb_i)e^{i\Phi} \tag{5.40}$$

(owing to (5.11) and (3.18)) and

$$F_{pq} = i(\rho^n_pk_nA_{\{k\}q}-\rho^n_qk_nA_{\{k\}p}) = [(\rho^0_p-\rho^1_p)A_{\{k\}q}-(\rho^0_q-\rho^1_q)A_{\{k\}p}]k_0i\,\mathrm{e}^{i\Phi} \tag{5.41}$$

(the transformation (3.15) has been used) with the components

$$F_{01} = \left[\frac{g}{2}-\gamma(k-g)+(\gamma h-\frac{g}{2}f)n^1\right]n^3\frac{(j\varkappa)^2}{L}b_3k_0i\,\mathrm{e}^{i\Phi}, \tag{5.42}$$

$$F_{02}=(\gamma h-\frac{g}{2}f)n^2n^3\frac{(j\varkappa)^2}{L}b_3k_0i\,\mathrm{e}^{i\Phi}, \tag{5.43}$$

$$F_{03}=(\gamma h-\frac{g}{2}f)n^3n^3\frac{(j\varkappa)^2}{L}b_3k_0i\,\mathrm{e}^{i\Phi}, \tag{5.44}$$

$$F_{12}=(\gamma-\frac{g}{2}k)(n^1-n^2)n^3\frac{(j\varkappa)^2}{L}b_3k_0i\,\mathrm{e}^{i\Phi}, \tag{5.45}$$

$$F_{13}=\Big[fk-h+(f-kh)n^1-(\gamma-\frac{g}{2}k)n^1n^1\Big]n^3n^3\frac{(j\varkappa)^2}{L}b_3k_0i\,\mathrm{e}^{i\Phi}, \tag{5.46}$$

$$F_{23}=\Big[f-kh-(\gamma-\frac{g}{2}k)n^1\Big]n^2\frac{(j\varkappa)^2}{L}b_3k_0i\,\mathrm{e}^{i\Phi}\,. \tag{5.47}$$

When

$$R^0=0, \tag{5.48}$$

the above representations are reduced to read merely

$$F_{01}\big|_{R^0=0}=\Big[\frac{g}{2}+\gamma g+(\gamma h+\frac{g^2}{4})n^1\Big]n^3|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}, \tag{5.49}$$

$$F_{02}\big|_{R^0=0}=(\gamma h+\frac{g^2}{4})n^2n^3|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}, \tag{5.50}$$

$$F_{03}\big|_{R^0=0}=(\gamma h+\frac{g^2}{4})n^3n^3|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}, \tag{5.51}$$

$$F_{12}\big|_{R^0=0}=\gamma(n^1-n^2)n^3|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}, \tag{5.52}$$

$$F_{13}\big|_{R^0=0}=(-h-\frac{g}{2}n^1-\gamma n^1n^1)n^3n^3|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}, \tag{5.53}$$

$$F_{23}\big|_{R^0=0}=(-\frac{g}{2}-\gamma n^1)n^2|\mathbf{R}|^{\frac{1}{2}(h-1)}Cb_3\,\mathrm{e}^{i\Phi_0}\,. \tag{5.54}$$

NOTE. Upon substituting (B.42) the Lagrangian density (3.4) takes on the form

$$\mathcal{D}^{pq}(R)=\sqrt{|\det(s_{pq}(g;R))|}\,s^{pt}(g;R)s^{qs}(g;R)F_{ts}(R) \tag{5.55}$$

that corresponds to the flat-space case, for the curvature tensor constructed from the conformal metric tensor $\{s_{pq}\}$ vanishes identically. This is the reason why the above Proposition 4 is valid. The respective transformation for the electromagnetic vector potential is

$$A_p(R)=\rho_p^m(g;r)B_m(r) \tag{5.56}$$

with

$$\rho_p^m=t_p^ik_i^m \tag{5.57}$$

(cf. (3.33) and (5.1)).

At the $O_1(g)$–level of consideration, the electromagnetic components (5.15)–(5.18) reduce to read

$$A_{\{k\}0} = \frac{1}{2} g \frac{|\mathbf{R}| R^3}{S^2} b_3 \, \mathrm{e}^{ik_p R^p}, \tag{5.58}$$

$$A_{\{k\}1} = -\frac{1}{2} g \frac{R^0}{|\mathbf{R}|} \frac{R^1 R^3}{S^2} b_3 \, \mathrm{e}^{ik_p R^p}, \qquad A_{\{k\}2} = -\frac{1}{2} g \frac{R^0}{|\mathbf{R}|} \frac{R^2 R^3}{S^2} b_3 \, \mathrm{e}^{ik_p R^p}, \tag{5.59}$$

$$A_{\{k\}3} = \left\{ 1 - \frac{1}{2} g \left[i |\mathbf{R}| k_0 + \frac{1}{2} (1 + i k_a R^a) \ln \frac{R^0 - |\mathbf{R}|}{R^0 + |\mathbf{R}|} + \frac{R^0}{|\mathbf{R}|} \frac{(R^3)^2}{S^2} \right] \right\} b_3 \, \mathrm{e}^{ik_p R^p}, \tag{5.60}$$

where $S^2 = (R^0)^2 - |\mathbf{R}|^2$ and (B.56) has been applied.

The *spherical waves of photons* are obtained upon taking the electromagnetic vector potential in the conformally flat space in the ordinary way:

$$B_{\{\omega j m\}0} = 0 \tag{5.61}$$

and

$$B_{\{\omega j m\}a}(r^q) = e^{ik_0 r^0} \int B_{\{\omega j m\}a}(\mathbf{k}) e^{i k_b r^b} \frac{d^3 k}{(2\pi)^3} \tag{5.62}$$

with

$$B_{\{\omega j m\}a}(\mathbf{k}) = \frac{4\pi^2}{\omega^{3/2}} \delta(|\mathbf{k}| - \omega) Y_{\{jm\}a}(\mathbf{k}/\omega), \tag{5.63}$$

where $\omega = k_0$ and $Y_{\{jm\}a}$ are required spherical functions subjected to the condition $\big(\mathbf{k}\mathbf{Y}_{\{jm\}}(\mathbf{k}/\omega)\big) = 0$. The quasi-pseudoeuclidean version reads

$$U_{\{\omega j m\}i}(t) = k_i^a(g;t) B_{\{\omega j m\}a}(r^q(g;t)) \tag{5.64}$$

and returning back into the $\mathcal{E}_g^{SR}$–space yields the components

$$A_{\{\omega j m\}p}(R) = \rho_p^a(g;R) B_{\{\omega j m\}a}(r^q(g;R)). \tag{5.65}$$

Since in the pseudoeuclidean approach proper the spherical electromagnetic waves are expanding with the light velocity value, which relates to $(r^0)^2 - \mathbf{r}^2 = 0$, then in the initial $\mathcal{E}_g^{SR}$–space we must respectively have $F(g;R) = 0$, whence we may be certain that the respective velocity value in the $\mathcal{E}_g^{SR}$–space is exactly $v = g_+$ (just in conformity with the velocity value (C.7) obtainable in case of the planar-type waves).

The *$\mathcal{E}_g^{SR}$–space extension of the Coulomb law* is given by the formulae:

$$B_0 = \frac{e}{\sqrt{\delta_{ab} r^a r^b}} = \frac{eh}{\xi(g;t)|\mathbf{t}|} = \frac{e}{j(g;R)\varkappa(g;R)|\mathbf{R}|} \tag{5.66}$$

(the formula (B.38) has been used in the second step and the formula (B.2) has been applied in the last step); in the quasi-pseudoeuclidean space,

$$U_i = k_i^0 B_0 \tag{5.67}$$

(see (B.38) and (B.39)) with the components

$$U_0 = \left(1 + (h-1)\frac{t^0 t_0}{S^2}\right) \frac{e}{|\mathbf{t}|}, \qquad U_a = (h-1)\frac{t^0 t_a}{S^2} \frac{e}{|\mathbf{t}|}; \tag{5.68}$$

in the $\mathcal{E}_g^{SR}$–space,

$$A_p = \rho_p^0 B_0 \tag{5.69}$$

(see (B.56)) with the components

$$A_0 = \left(1 + \frac{1}{L}(h - 1 + \frac{g}{2}\frac{R^0}{|\mathbf{R}|} - \frac{g^2}{4})\right)\frac{e}{|\mathbf{R}|}, \qquad A_a = -\left((h-1)\frac{R^0}{|\mathbf{R}|} + \frac{g}{2}\right)\frac{R^a e}{|\mathbf{R}|^2}. \tag{5.70}$$

In the $O_1(g)$–approximation we get merely

$$U_0 = \frac{e}{|\mathbf{t}|}, \qquad U_a = 0, \tag{5.71}$$

and

$$A_0 = \frac{e}{|\mathbf{R}|} + \frac{g}{2}\frac{R^0 e}{S^2}, \qquad A_a = -\frac{g}{2}\frac{R^a e}{|\mathbf{R}|^2}. \tag{5.72}$$

6. Conformal scalar and spinor fields and solutions

In distinction from the electromagnetic field case, the conventional Lagrangian (2.2) for a complex scalar field $\phi = \phi(R)$ is not conformally invariant. Let us modify the Lagrangian as follows:

$$L_{\{\phi\}} = g^{pq}(g;R)\partial_p\phi^*\partial_q\phi + \frac{1}{6}S^{pq}{}_{pq}\phi^*\phi - \varkappa^2 m^2\phi^*\phi. \tag{6.1}$$

Since

$$S^{pq}{}_{pq} = \frac{1}{4}g^2\frac{6}{F^2(g;R)} \tag{6.2}$$

(see (A.22)–(A.23)), we get

$$L_{\{\phi\}} = g^{pq}(g;R)\partial_p\phi^*\partial_q\phi + \frac{1}{4}g^2\frac{1}{F^2}\phi^*\phi - \varkappa^2 m^2\phi^*\phi. \tag{6.3}$$

This Lagrangian entails the associated scalar field equation

$$\frac{1}{J}\frac{\partial}{\partial R^p}\left(Jg^{pq}\frac{\partial}{\partial R^q}\phi\right) - \frac{1}{4}g^2\frac{1}{F^2}\phi + \varkappa^2 m^2\phi = 0 \tag{6.4}$$

which extends (2.8)–(2.9). Using the coordinates

$$r^n = \rho^n(g;R), \tag{6.5}$$

where ρ^n are the functions given by (B.44), let us go over to a new field

$$a = a(r) \tag{6.6}$$

according to the transformation

$$\phi(R) = \varkappa(g;R)a(\rho(g;R)) \tag{6.7}$$

($\varkappa$ is the conformal multiplier (B.43)). Calculations show that the insertion of (6.7) into (6.4) results in the equation of the ordinary Klein-Gordon type:

$$e^{ij}\frac{\partial^2 a}{\partial r^i \partial r^j} + m^2 a = 0 \tag{6.8}$$

which is obviously derivable from the ordinary pseudoeuclidean Lagrangian

$$L_{\{a\}} = e^{ij}\frac{\partial a^*}{\partial r^i}\frac{\partial a}{\partial r^j} - m^2 a^* a. \tag{6.9}$$

Accordingly, we introduce

DEFINITION. The field $a = a(r)$ is called the *conformal scalar field.*

In particular, the plane-wave solutions to the latter field generate the solutions

$$\phi(R) = \frac{1}{(2\pi)^{3/2}}\varkappa(g;R)\int \mathrm{e}^{i\Phi}\,\delta(k^2 - m^2)\phi(k)d^4k, \tag{6.10}$$

where Φ is the phase (B.55). By using (2.19), the latter solutions can be transformed to the quasi-pseudoeuclidean space:

$$u(t) = \frac{1}{(2\pi)^{3/2}}\xi(g;t)\int \mathrm{e}^{i\Phi}\,\delta(k^2 - m^2)u(k)d^4k, \tag{6.11}$$

in which case the representation (B.51) for the phase Φ should be taken.

The quasi-pseudoeuclidean versions of the formulae (6.3), (6.4), and (6.7) read

$$L_{\{u\}} = n^{ij}(g;t)\partial_i u^* \partial_j u + \frac{1}{4}g^2\frac{1}{S^2}u^* u - \xi^2 m^2 u^* u, \tag{6.12}$$

$$\partial_i(n^{ij}u_j) - \frac{1}{4}g^2\frac{1}{S^2}u + \xi^2 m^2 u = 0, \tag{6.13}$$

and

$$u(t) = \xi(g;t)a\Big(\frac{1}{h}\xi(g;t)t\Big), \tag{6.14}$$

where $u = u(t) = \phi(R)$ in accordance with (2.19) and ξ is the conformal multiplier (B.36). We get

$$u_i = \frac{\gamma t_i}{S^2}\xi a + \frac{1}{h}\xi^2 a_i + \frac{1}{h}\gamma\xi^2\frac{a_n t^n}{S^2}t_i, \qquad t^i u_i = \gamma\xi a + \xi^2 a_n t^n, \tag{6.15}$$

with $a_n(r) = \dfrac{\partial a(r)}{\partial r^n}$ and $u_i = \dfrac{\partial u}{\partial t^i}$. Also,

$$n^{ij}u_j = \gamma\xi a\frac{t^i}{S^2} - \gamma\xi^2\frac{a_n t^n}{S^2}t_i + k\xi^2 a^i, \tag{6.16}$$

where $a^i = e^{ij}a_j$,

$$\partial_j a_i = \frac{1}{h}\xi(a_{ij} + \gamma\frac{a_{ik}t^k}{S^2}t_j), \tag{6.17}$$

and

$$a_{ij}(r) = \frac{\partial^2 a(r)}{\partial r^i \partial r^j}. \tag{6.18}$$

Finally

$$\partial_i(n^{ij}u_j) = \frac{g^2}{4}\xi\frac{a}{S^2} + \xi^3\frac{\partial a^i}{\partial r^i}. \tag{6.19}$$

Inserting the last result in (6.13) just yields (6.8).

The $\mathcal{E}_g^{SR}$–*space extension of the Yukawa potential* is given by the following formulae:

$$a=\frac{\exp\Bigl(-m\sqrt{\delta_{ab}r^ar^b}\Bigr)}{m\sqrt{\delta_{ab}r^ar^b}}=\frac{h\exp\Bigl(-\frac{1}{h}m\xi(g;t)|\mathbf{t}|\Bigr)}{m\xi(g;t)|\mathbf{t}|}=\frac{\exp\Bigl(-mj(g;R)\varkappa(g;R)|\mathbf{R}|\Bigr)}{mj(g;R)\varkappa(g;R)|\mathbf{R}|}, \tag{6.20}$$

$$u(t)=\frac{h}{m|\mathbf{t}|}\exp\Bigl(-\frac{1}{h}m\xi(g;t)|\mathbf{t}|\Bigr) \tag{6.21}$$

(in accordance with the rule (6.14)), and

$$\phi(R)=\frac{1}{mj(g;R)|\mathbf{R}|}\exp\Bigl(-mj(g;R)\varkappa(g;R)|\mathbf{R}|\Bigr). \tag{6.22}$$

In the $O_1(g)$–approximation these representations reduce to

$$u(t)=\frac{1}{m|\mathbf{t}|}\,\mathrm{e}^{-m|\mathbf{t}|} \tag{6.23}$$

and

$$\phi(R)=\frac{1}{m|\mathbf{R}|}\left(1+\frac{1}{4}g(1+m|\mathbf{R}|)\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|}\right)\mathrm{e}^{-m|\mathbf{R}|}\,. \tag{6.24}$$

It is worth comparing (6.20)–(6.24) with (5.61)–(5.67).

For the spinor we take the Lagrangian

$$L_{\{\psi\}}=\frac{i}{2}[\bar\psi\gamma^pD_p\psi-(D_p\bar\psi)\gamma^p\psi]-\varkappa m\bar\psi\psi \tag{6.25}$$

in which the factor $\varkappa$ in the last term is the only distinction from the standard choice. This entails the following generalization of the Dirac equation:

$$i\gamma^pD_p\psi-\varkappa m\psi=0. \tag{6.26}$$

Applying the substitution

$$\psi(R)=[\varkappa(g;R)]^{\alpha}U(\rho(g;R)) \tag{6.27}$$

with

$$\alpha=\frac{3}{2} \tag{6.28}$$

reduces the equation (6.22) to the ordinary pseudoeuclidean form

$$i\widetilde\gamma^iD_iU-mU=0, \tag{6.29}$$

which can be verified by straightforward calculations; the notation $\widetilde\gamma^i=\gamma^Ph_P^i$ has been used. So,

$$L_{\{U\}}=\frac{i}{2}[\bar U\widetilde\gamma^iD_iU-(D_i\bar U)\widetilde\gamma^iU]-m\bar UU. \tag{6.30}$$

Accordingly, we introduce

DEFINITION. The field $U=U(r)$ is called the *conformal spinor field.*

In particular, the equation (6.22) admits the wave solution

$$\psi(R)=\frac{1}{(2\pi)^{3/2}}[\varkappa(g;R)]^{\alpha}\int \mathrm{e}^{i\Phi}\,\delta(k^2-m^2)\psi(k)d^4k, \tag{6.31}$$

where, as in the ordinary theory, the amplitude $\psi(k)$ is subjected by the condition

$$(\gamma^P k_P+m)\psi(k)\Big|_{k^2=m^2}=0. \tag{6.32}$$

By means of the transformation (4.11) the solution can be transformed to the quasi-pseudoeuclidean space:

$$v(t)=\frac{1}{(2\pi)^{3/2}}[\xi(g;t)]^{\alpha}\int \mathrm{e}^{i\Phi}\,\delta(k^2-m^2)v(k)d^4k, \tag{6.33}$$

where $v(k)=\psi(k)$.

The quasi-pseudoeuclidean versions of the formulae (6.21)–(6.23) read

$$L_{\{v\}}=\frac{i}{2}[\bar{v}\gamma^i D_i v-(D_i\bar{v})\gamma^i v]-\xi m\bar{v}v, \tag{6.34}$$

$$i\gamma^i D_i v-\xi m v=0, \tag{6.35}$$

and

$$v(t)=[\xi(g;t)]^{\alpha}U(\frac{1}{h}\xi(g;t)t). \tag{6.36}$$

We find

$$v_n(t)=\gamma\alpha\frac{t_n}{S^2}\xi^{\alpha}U+\xi^{\alpha+1}f_n^m U_m, \tag{6.37}$$

$$\frac{\partial U}{\partial t^n}=\xi\frac{1}{h}(U_n+\frac{\gamma}{S^2}t_n t^m U_m)=\xi f_n^m U_m, \tag{6.38}$$

where $v_n(t)=\partial v(t)/\partial t^n$ and $U_n(r)=\partial U(r)/\partial r^n$, and also

$$\gamma^n v_n=\gamma\alpha\frac{\gamma_j t^j}{S^2}\xi^{\alpha}U+\xi^{\alpha+1}\tilde{\gamma}^m U_m \tag{6.39}$$

together with

$$\gamma^n D_n v=\gamma^n(\partial_n v+\frac{1}{8}R^{PQ}{}_n(\gamma_P\gamma_Q-\gamma_Q\gamma_P)v)=\gamma(\alpha-\frac{3}{2})\frac{\gamma_j t^j}{S^2}\xi^{\alpha}U+\xi^{\alpha+1}\tilde{\gamma}^m U_m \tag{6.40}$$

(the formulas (4.12) and (4.13) together with (B.35) have been used). Comparing the last result with (6.24) and (6.31) verifies (6.25).

7. $\mathcal{E}_g^{SR}$–space quantization of fields

Starting from the conformally flat space referred to the coordinates r^n given by (6.5), the results of the ordinary relativistic quantum fields can be used.

We obtain the following wave decompositions: for the scalar field

$$\phi(R)=\frac{1}{(2\pi)^{3/2}}\varkappa(g;R)\int \mathrm{e}^{ik_n r^n(g;R)}\,\phi(k)\delta(k^2-m^2)d^4k, \tag{7.1}$$

for the electromagnetic vector potential

$$A_p(R) = \frac{1}{(2\pi)^{3/2}}\rho_p^j(g;R)\int a_j(k_n)\,\mathrm{e}^{ik_n r^n(g;R)}\,\delta(k^2)d^4k, \tag{7.2}$$

and for the spinor field

$$\psi(R) = \frac{1}{(2\pi)^{3/2}}[\varkappa(g;R)]^{3/2}\int \mathrm{e}^{ik_n r^n(g;R)}\,\psi(k)\delta(k^2-m^2)d^4k. \tag{7.3}$$

The waves obey the proper value equations in single-wave cases:

$$\mathcal{P}_n\phi_{\{k\}} = k_n\phi_{\{k\}}, \qquad \mathcal{P}_nA_{\{k\}q} = k_nA_{\{k\}q}, \qquad \mathcal{P}_n\psi_{\{k\}} = k_n\psi_{\{k\}}, \tag{7.4}$$

where $\mathcal{P}_n$ is the *operator of the four-dimensional momentum* which acts in the following way:

$$\mathcal{P}_n\phi_{\{k\}} = -i\eta_n^q\partial_q(\frac{1}{\varkappa}\phi_{\{k\}}) \tag{7.5}$$

in scalar case,

$$\mathcal{P}_n\psi_{\{k\}} = -i\eta_n^q\partial_q(\frac{1}{\varkappa^{3/2}}\psi_{\{k\}}) \tag{7.6}$$

in spinor case, and simply

$$\mathcal{P}_nA_q = -i\eta_n^q(\partial_qA_p - \eta_m^s\partial_q\rho_p^mA_s) \tag{7.7}$$

in electromagnetic case; the coefficients ρ_p^m and η_n^q are given by (B.48) and (B.49).

The generalized differences

$$(t\ominus t')^n = \frac{1}{h}\xi(g;t)t^n - \frac{1}{h}\xi(g;t')t'^n \tag{7.8}$$

(see (B.38)) and

$$(R\ominus R')^n = \rho^n(g,R) - \rho^n(g;R') \tag{7.9}$$

(see (B.44)) are arisen.

The quasi-pseudoeuclidean versions to the representations (7.1)–(7.7) read: for the scalar field

$$u(t) = \frac{1}{(2\pi)^{3/2}}\xi(g;t)\int \mathrm{e}^{i\xi(g;t)k_nt^n/h}\,\phi(k)\delta(k^2-m^2)d^4k, \tag{7.10}$$

for the electromagnetic vector potential

$$U_i(t) = \frac{1}{(2\pi)^{3/2}}k_i^j(g;t)\int a_j(k_n)\,\mathrm{e}^{i\xi(g;t)k_nt^n/h}\,\delta(k^2)d^4k, \tag{7.11}$$

and for the spinor field

$$v(t) = \frac{1}{(2\pi)^{3/2}}[\xi(g;t)]^{3/2}\int \mathrm{e}^{i\xi(g;t)k_nt^n/h}\,v(k)\delta(k^2-m^2)d^4k; \tag{7.12}$$

also,

$$\mathcal{P}_nu_{\{k\}} = k_nu_{\{k\}}, \qquad \mathcal{P}_nU_{\{k\}j} = k_nU_{\{k\}j}, \qquad \mathcal{P}_nv_{\{k\}} = k_nv_{\{k\}}, \tag{7.13}$$

with

$$\mathcal{P}_nu_{\{k\}} = -ik_n^j\partial_j(\frac{1}{\xi}u_{\{k\}}), \tag{7.14}$$

$$\mathcal{P}_n v\psi_{\{k\}} = -ik^j_n\partial_j(\frac{1}{\xi^{3/2}}v_{\{k\}}), \tag{7.15}$$

and

$$\mathcal{P}_nU_j = -ik^j_n(\partial_jU_i - k^l_m\partial_jK^m_iU_l), \tag{7.16}$$

where k^i_j are the coefficients (B.39) and $K^n_ik^i_j = \delta^n_j$.

In case of scalar field, the commutator of the operators $\phi^-(\mathbf{k})$ and $\phi^+(\mathbf{q})$ is subjected to the condition

$$[\phi^-(\mathbf{k}), \phi^+(\mathbf{q})] = \delta(\mathbf{k} - \mathbf{q}). \tag{7.17}$$

In case of the electromagnetic field, it is appropriate to take the decomposition

$$B_m = e^s_m b^\pm_s(\mathbf{k})\,\mathrm{e}^{i\phi} \tag{7.18}$$

in terms of an orthonormal frame $\{e^s_i\}$, so that $e^m_ie^k_je_{mk} = e_{ij}$, assume the property

$$b^+_i(\mathbf{k})b^{i,-}(\mathbf{k}) = e^{mn}a^+_m(\mathbf{k})a^-_n(\mathbf{k}), \tag{7.19}$$

and set forth the commutator condition

$$[a^-_i(\mathbf{k}), a^+_j(\mathbf{q})] = -e_{ij}\delta(\mathbf{k} - \mathbf{q}). \tag{7.20}$$

To quantize the Fermion field, we regard the amplitudes $a_s(\mathbf{k})$ as some operators which satisfy the permutation relations of Fermi-Dirac type:

$$\{(a^-_s(\mathbf{k}))^*, a^+_r(\mathbf{q})\} = \{a^-_s(\mathbf{k}), (a^+_r(\mathbf{q}))^*\} = \delta_{sr}\delta(\mathbf{k} - \mathbf{q}). \tag{7.21}$$

So, the required commutators for fields in the $\mathcal{E}^{SR}_g$–space under consideration read as follows: in scalar case,

$$[\phi^-(R), \phi^+(R')] = \frac{1}{i}\varkappa(g;R)\varkappa(g;R')\Delta^-(R\ominus R'), \tag{7.22}$$

$$[\phi^+(R), \phi^-(R')] = \varkappa(g;R)\varkappa(g;R')\frac{1}{i}\Delta^+(R\ominus R'), \tag{7.23}$$

and

$$[\phi(R), \phi(R')] = \varkappa(g;R)\varkappa(g;R')\frac{1}{i}\Delta(R\ominus R'); \tag{7.24}$$

in electromagnetic case,

$$[A^-_p(R), A^+_q(R')] = is_{pq}(g;R,R')\Delta^-_0(R\ominus R'), \tag{7.25}$$

$$[A^+_p(R), A^-_q(R')] = is_{pq}(g;R,R')\Delta^+_0(R\ominus R'), \tag{7.26}$$

and

$$[A_p(R), A_q(R')] = is_{pq}(g;R,R')\Delta_0(R\ominus R'), \tag{7.27}$$

where the tensor

$$s_{pq}(g;R,R') = \rho^m_p(g;R)\rho^k_q(g;R')e_{mk} \tag{7.28}$$

appears; in spinor case,

$$[\psi(R), \bar{\psi}(R')] = \frac{1}{i}[\varkappa(g;R)\varkappa(g;R')]^{3/2}S(R \ominus R'). \tag{7.29}$$

Converting the consideration into the quasi-pseudoeuclidean space, we obtain the following result for the respective commutators: for the scalar field,

$$[u^-(t), u^+(t')] = \xi(g;t)\xi(g;t')\frac{1}{i}\Delta^-(t \ominus t'), \tag{7.30}$$

$$[u^+(t), u^-(t')] = \xi(g;t)\xi(g;t')\frac{1}{i}\Delta^+(t \ominus t'), \tag{7.31}$$

and

$$[u(t), u(t')] = \xi(g;t)\xi(g;t')\frac{1}{i}\Delta(t \ominus t'); \tag{7.32}$$

for the electromagnetic field

$$[U_i^-(t), U_j^+(t')] = ic_{ij}(g;t,t')\Delta_0^-(t \ominus t'), \tag{7.33}$$

$$[U_i^+(t), U_j^-(t')] = ic_{ij}(g;t,t')\Delta_0^+(t \ominus t'), \tag{7.34}$$

and

$$[U_i(t), U_j(t')] = ic_{ij}(g;t,t')\Delta_0(t \ominus t'), \tag{7.35}$$

where

$$c_{ij}(g;t,t') = k_i^m(g;t)k_j^k(g;t')e_{mk}; \tag{7.36}$$

for the spinor field

$$[v(t), \bar{v}(t')] = \frac{1}{i}[\xi(g;t)\xi(g;t')]^{3/2}S(t \ominus t'). \tag{7.37}$$

In the very conformally flat space, the ordinary pseudoeuclidean relations take place, so that in scalar case,

$$[\phi^-(r), \phi^+(r')] = \frac{1}{i}\Delta^-(r - r'),$$

$$[\phi^+(r), \phi^-(r')] = \frac{1}{i}\Delta^+(r - r'),$$

$$[\phi(r), \phi(r')] = \frac{1}{i}\Delta(r - r');$$

in electromagnetic case,

$$[B_i^-(r), B_j^+(r')] = ie_{ij}\Delta_0^-(r - r'),$$

$$[B_i^+(r), B_j^-(r')] = ie_{ij}\Delta_0^+(r - r'),$$

$$[B_i(r), B_j(r')] = ie_{ij}\Delta_0(r - r');$$

and in spinor case,

$$[U(r),\bar{U}(r')]=\frac{1}{i}S(r-r');$$

with the permutation functions

$$\Delta^{+}(r)=\frac{1}{(2\pi)^3 i}\int\theta(k_0)e^{ik_n r^n}\delta(k^2-m^2)d^4k,$$

$$\Delta^{-}(r)=\frac{i}{(2\pi)^3}\int\theta(-k_0)e^{-ik_n r^n}\delta(k^2-m^2)d^4k=-\Delta^{+}(r),$$

$$\Delta(r)=\Delta^{+}(r)+\Delta^{-}(r)=\frac{i}{(2\pi)^3}\int\epsilon(k_0)e^{-ik_n r^n}\delta(k^2-m^2)d^4k,$$

$$\Delta_0^{+}(r)=\frac{1}{(2\pi)^3 i}\int\theta(k_0)e^{ik_n r^n}\delta(k^2)d^4k,$$

$$\Delta_0^{-}(r)=\frac{i}{(2\pi)^3}\int\theta(-k_0)e^{-ik_n r^n}\delta(k^2)d^4k=-\Delta_0^{+}(r),$$

$$\Delta_0(r)=\Delta_0^{+}(r)+\Delta_0^{-}(r)=\frac{i}{(2\pi)^3}\int\epsilon(k_0)e^{-ik_n r^n}\delta(k^2)d^4k,$$

and

$$S(r)=(i\gamma^j\frac{\partial}{\partial r^j}+m)\Delta(r).$$

8. $\hat{\mathcal{E}}_g^{SR}$–way regularization

The presence of the geometrical parameter g can be used for constructing an appropriate function $\mathcal{W}$ which ought to play the role of the $\hat{\mathcal{E}}_g^{SR}$*–space weight* when performing integration over the four-dimensional momenta $\{P_q\}\in\hat{V}_4$.

Suppressing ultraviolet divergences can be motivated, first of all, by the circumstance that, since the $\hat{\mathcal{E}}_g^{SR}$–space approach under study is based upon a group of invariance (which transformations leave the FHF H in the co-space $\hat{V}_4$ invariant), the function $\mathcal{W}$ should include the dependence on $\{P_q\}$ through the modulus $P=H(g;P_q)$. Next, when integrating over $\{P_q\}$, it is natural to choose the function $\mathcal{W}$ to be a decreasing exponent:

$$\mathcal{W}(g;P)=C_1(g)e^{-\alpha(g)P^{\nu}}, \tag{8.1}$$

where $\alpha(g)>0$, $\nu>0$, and $C_1(g)$ stands for a positive normalizing factor, so that

$$\int_{P=0}^{\infty}\mathcal{W}(g;P)P^3dP=1 \tag{8.2}$$

and

$$C_1(g)=\frac{1}{\int_{P=0}^{\infty}\mathcal{W}(g;P)P^3dP}. \tag{8.3}$$

The answer to the question “what choice is to be preferred for the factor $\alpha(g)$” can tentatively be given by realizing the attractive idea of geometrical origin that the factor stems from the $\hat{\mathcal{E}}_g^{SR}$–space curvature. For the later is proportional to g^2 (see (A.22) and (A.23)), we suggest the choice

$$\alpha = Cg^2, \tag{8.4}$$

where C is a positive constant. The respective invariant integration with the $\hat{\mathcal{E}}_g^{SR}$–space weight $\mathcal{W}$ over the space $\hat{V}_4$ should be performed in accordance with the rule

$$\int_{\hat{V}_4} \mathcal{M} = \int_{P=0}^{\infty} Z(g;P)\mathcal{W}(g;P)P^3 dP, \qquad Z = \int \mathcal{M}\hat{J}d\hat{\Omega}, \tag{8.5}$$

where $\mathcal{M}$ is some matrix element or scalar, $\hat{J} = \sqrt{|\det(g_{pq}(g;P_s))|}$, and $d\hat{\Omega}$ is an appropriate angular element of volume. Because of the exponential form (8.1) of the weight $\mathcal{W}$, the integral with respect to P will be converged at the higher limit for any degrees z of the terms proportional to P^z. Therefore, upon the Finslerian extension under consideration, the finite expressions proportional to natural degrees of the square of the inversed $\hat{\mathcal{E}}_g^{SR}$–space parameter, $1/g^2$, will relate to the divergent integrals that appear in ordinary theory at $P \to \infty$.In this sense the $\hat{\mathcal{E}}_g^{SR}$–space approach developed may give, in principle, a simple explanation for the nature of the ultraviolet divergences (which is characteristic for the ordinary Lorentz-invariant theory of quantized fields:the limiting transition $g \to 0$ in the factor (8.3) is far from always being possible! Though in the exponent of the $\hat{\mathcal{E}}_g^{SR}$–space weight (8.1) it is possible to go over to the limit $g \to 0$, which does remove the exponent (for $e^{-Cg^2P^\nu} \to 1$ at $g \to 0$), the normalizing factor $C_1(g)$ cannot be analytical at $g = 0$. For example, it is worth choosing

$$\mathcal{W}(g;P) = C_1(g)e^{-g^2P^2/m^2}, \tag{8.6}$$

where $m \neq 0$ is the mass of particle, and use the Fourier transformation

$$\frac{1}{(2\pi)^4}\int \mathrm{e}^{ik_n r^n - g^2(k^2/m^2)}\, h^{-2} d^4k = \frac{1}{16\pi^2 h^2}\frac{m^4}{g^4}\,\mathrm{e}^{-r^2m^2/4g^2}\,. \tag{8.7}$$

In this way, the so-called alpha-representation

$$\frac{1}{m^2 - k^2 - i\epsilon} = i\int \mathrm{e}^{i\alpha(k^2 - m^2 + i\epsilon)}\, d\alpha \tag{8.8}$$

can be applied to find form-factors and evaluate various diagrams and matrix elements.

Suppressing angular divergences, which are stemming from integrations over the particle energy variable P_0, is of an alternative need. By following the known general ideas and methods of statistical physics, we can take the respective weight $\mathcal{W}_1$ in the form of the exponent

$$\mathcal{W}_1 = C_3(g)e^{-C_2g^2P_0}, \tag{8.9}$$

where C_2 and C_3 are positive constants and P_0 is the particle energy. To have in the exponent the energy in dimensionless combination, we may substitute the normalized energy

$$K_0 = P_0/m \tag{8.10}$$

with P_0. In the on-mass-shell treatment, that is, when one sets forth the dispersion relation

$$H(g;P_q) = m \tag{8.11}$$

(with the FHF H given by (A.14)), the phase volume element is to be taken as

$$d\hat{V} \overset{\text{def}}{=} \delta\left((H(g;P_q))^2 - m^2\right)\hat{J}d^4P = \hat{J}\frac{d^3\mathbf{P}}{2P^0}\Big|_{H(g;P_q)=m} \tag{8.12}$$

and the weight takes on the form

$$\mathcal{W}_1(g,|\mathbf{P}|) = C_3(g)e^{-C_4g^2K_0(g;|\mathbf{P}|/m)} \tag{8.13}$$

with the function $K_0 = K_0(g;|\mathbf{P}|/m)$ obtainable from the equation $H(g;K_0,\mathbf{P}/m) = 1$. Because of the exponential form (8.13) the integral over $|\mathbf{P}|$ will converge at the higher limit at any degrees $|\mathbf{P}|^n$ in the matrix element, so that the divergent integrals appeared in the ordinary pseudo-euclidean theory at the process $|\mathbf{P}| \to \infty$,upon $\mathcal{E}_g^{SR}$–space extension can be juxtaposed with finite expressions proportional to degrees of $1/g^2$. On neglecting the parameter g in the function $K_0 = K_0(g;|\mathbf{P}|/m)$, the weight function (8.12) is simplified to read

$$\mathcal{W}_1(g,|\mathbf{P}|) = C_3(g)e^{-C_4g^2\sqrt{1+(|\mathbf{P}|/m)^2}}. \tag{8.14}$$

In such a case, the normalization

$$\int \mathcal{W}_1(g,|\mathbf{P}|)d^3\mathbf{P} = 1 \tag{8.15}$$

entails

$$\frac{1}{4\pi C_3(g)} = 2\left(\frac{8}{g^2}\right)^2 K_1\left(\frac{g^2}{8}\right) + \left(\frac{8}{g^2}\right)K_0\left(\frac{g^2}{8}\right), \tag{8.16}$$

where $\acute{K}_0$ and $\acute{K}_1$ — the Macdonald functions (the Hankel functions of imaginary argument), for which

$$\int_0^\infty e^{-z\,\mathrm{ch}\,u}\,\mathrm{sh}^2\,u\,du = \frac{1}{z}\acute{K}_1(z) \tag{8.17}$$

and

$$\acute{K}_1'(z) = -\frac{1}{z}\acute{K}_1(z) - \acute{K}_0(z). \tag{8.18}$$

In the law-energy region

$$\frac{|\mathbf{P}|}{m} \ll 1 \tag{8.19}$$

we get the Maxwell-type distribution

$$\mathcal{W}_1(g,|\mathbf{P}|) = C_3(g)e^{-C_4g^2|\mathbf{P}|^2/2m^2}. \tag{8.20}$$

Alternatively, in the ultrarelativistic region

$$\frac{|\mathbf{P}|}{m} \gg 1 \tag{8.21}$$

we get the approximation

$$\mathcal{W}_1(g,|\mathbf{P}|) = C_3(g)e^{-C_4g^2|\mathbf{P}|/m}. \tag{8.22}$$

Appendix A. Basic properties of the space $\mathcal{E}_g^{SR}$

Searching for extension of the pseudoeuclidean geometry in due Finsler-relativistic way, we should adapt constructions to the following decomposition of the four-dimensional vector space V_4:

$$V_4 = \mathcal{S}_g^+ \cup \Sigma_g^+ \cup \mathcal{R}_g \cup \Sigma_g^- \cup \mathcal{S}_g^-, \tag{A.1}$$

which sectors relate to the cases when the contravariant vector $R \in V_4$ is respectively future–timelike, future–isotropic, spacelike, past–isotropic, and past–timelike. The respective co-analogue for the covariant vectors (the momenta) $P \in \hat{V}_4$ reads

$$\hat{V}_4 = \hat{\mathcal{S}}_g^+ \cup \hat{\Sigma}_g^+ \cup \hat{\mathcal{R}}_g \cup \hat{\Sigma}_g^- \cup \hat{\mathcal{S}}_g^-. \tag{A.2}$$

With this purpose, we introduce the following convenient notation:

$$G = g/h, \tag{A.3}$$

$$h = \sqrt{1 + \frac{1}{4}g^2}, \tag{A.4}$$

$$g_+ = -\frac{1}{2}g + h, \qquad g_- = -\frac{1}{2}g - h, \tag{A.5}$$

$$G_+ = g_+/h \equiv -\frac{1}{2}G + 1, \qquad G_- = g_-/h \equiv -\frac{1}{2}G - 1, \tag{A.6}$$

$$g^+ = 1/g_+ = -g_-, \qquad g^- = 1/g_- = -g_+, \tag{A.7}$$

$$g^+ = \frac{1}{2}g + h, \qquad g^- = \frac{1}{2}g - h, \tag{A.8}$$

$$G^+ = g^+/h \equiv \frac{1}{2}G + 1, \qquad G^- = g^-/h \equiv \frac{1}{2}G - 1. \tag{A.9}$$

We shall decompose vectors to select the timelike components and the three-dimensional spatial components: $R = \{R^0, \mathbf{R}\}$ and $P = \{P_0, \mathbf{P}\}$. In terms of the forms

$$B(g;R) = -\left(R^0 + g_-|\mathbf{R}|\right)\left(R^0 + g_+|\mathbf{R}|\right) \equiv -\left((R^0)^2 - gR^0|\mathbf{R}| - |\mathbf{R}|^2\right) \tag{A.10}$$

and

$$\hat{B}(g;P) = -\left(P_0 - \frac{|\mathbf{P}|}{g^+}\right)\left(P_0 - \frac{|\mathbf{P}|}{g^-}\right) \equiv -\left((P_0)^2 + gP_0|\mathbf{P}| - |\mathbf{P}|^2\right), \tag{A.11}$$

all the sectors that enter the decompositions (A.1) and (A.2) can be embraced by one FMF

$$F(g;R) = \sqrt{|B(g;R)|}\, j(g;R) = \left|R^0 + g_-|\mathbf{R}|\right|^{G_+/2} \left|R^0 + g_+|\mathbf{R}|\right|^{-G_-/2}, \tag{A.12}$$

where

$$j(g;R) = \left|\frac{R^0 + g_-|\mathbf{R}|}{R^0 + g_+|\mathbf{R}|}\right|^{-G/4}, \tag{A.13}$$

and one FHF

$$H(g;P)=\sqrt{|\hat{B}(g;P)|}\,\hat{j}(g;P)=\left|P_0-\frac{|\mathbf{P}|}{g^+}\right|^{G^+/2}\left|P_0-\frac{|\mathbf{P}|}{g^-}\right|^{-G^-/2}, \tag{A.14}$$

where

$$\hat{j}(g;P)=\left|\frac{P_0-\dfrac{|\mathbf{P}|}{g^+}}{P_0-\dfrac{|\mathbf{P}|}{g^-}}\right|^{G/4}. \tag{A.15}$$

By following the methods of the Finsler geometry, we use the definitions for the covariant vector

$$R_p\overset{\text{def}}{=}\frac{1}{2}\frac{\partial F^2(g;R)}{\partial R^p}\equiv P_p$$

and the FMT

$$g_{pq}(g;R)\overset{\text{def}}{=}\frac{1}{2}\frac{\partial^2F^2(g;R)}{\partial R^p\partial R^q}=\frac{\partial R_p(g;R)}{\partial R^q};$$

$p,q,...=0,1,2,3.$

Thus we get the *pseudo-Finsleroid relativistic space*

$$\mathcal{E}_g^{SR}=\{V_4;\ F(g;R);\ g_{pq}(g;R);\ R\in V_4\} \tag{A.16}$$

and the *pseudo-Finsleroid relativistic co-space*

$$\hat{\mathcal{E}}_g^{SR}=\{\hat{V}_4;\ H(g;P);\ g^{pq}(g;P);\ P\in\hat{V}_4\}. \tag{A.17}$$

Special calculations can be used to verify the equalities

$$\det(g_{pq}(g;R))=-[j(g;R)]^8 \tag{A.18}$$

and

$$\operatorname{sign}(g_{pq})=\operatorname{sign}(g^{pq})=(+---). \tag{A.19}$$

The following assertion is valid: *for the pseudo-Finsleroid space $\mathcal{E}_g^{SR}$ the Cartan torsion tensor*

$$C_{pqr}\overset{\text{def}}{=}\frac{1}{2}\frac{\partial g_{pq}}{\partial R^r}$$

is of the special algebraic form

$$C_{pqr}=\frac{1}{N}\left(h_{pq}C_r+h_{pr}C_q+h_{qr}C_p-\frac{1}{C_tC^t}C_pC_qC_r\right), \tag{A.20}$$

where

$$C_pC^p=-\frac{N^2g^2}{4F^2} \tag{A.21}$$

and $N=4$. Here, $C_p=C_p{}^q{}_q$.

Proof is gained by straightforward calculations on the basis of the explicit form of components of the FMT and the Cartan tensor (see more detail in [9–11]). Inserting (A.20)–(A.21) in the general expression for the curvature tensor

$$S_{pqrs}=C_{tqr}C_p{}^t{}_s-C_{tqs}C_p{}^t{}_r$$

yields the following simple result after rather simple straightforward calculations:

$$S_{pqrs} = S^*(h_{pr}h_{qs} - h_{ps}h_{qr})/F^2 \tag{A.22}$$

with the constant

$$S^* = \frac{1}{4}g^2. \tag{A.23}$$

The tensor

$$h_{pr} \overset{\text{def}}{=} g_{pr} - l_p l_r \tag{A.24}$$

has been used, where $l_p = R_p/F(g;R)$ — the unit vector components.

The FMF (A.12) defines the *pseudo-Finsleroid*

$$\mathcal{F}_g^{\text{Relativistic}} := \{R \in V_4 : F(g;R) \le 1\}. \tag{A.25}$$

The associated *indicatrix* $\mathcal{I}_g$ defined by

$$\mathcal{I}_g := \{R \in V_4 : F(g;R) = 1\} \tag{A.26}$$

is the surface of the pseudo-Finsleroid. With the given FHF (A.14), the body

$$\hat{\mathcal{F}}_g^{\text{Relativistic}} := \{\hat{R} \in \hat{V}_4 : H(g;\hat{R}) \le 1\} \tag{A.27}$$

is called the *co-pseudo-Finsleroid.* The respective figuratrix introduced according to

$$\hat{\mathcal{I}}_g := \{\hat{R} \in \hat{V}_4 : H(g;\hat{R}) = 1\} \tag{A.28}$$

is called the *co-indicatrix.*

From (A.22)–(A.24) it follows that *in case of the pseudo-Finsleroid space* $\mathcal{E}_g^{SR}$ *the indicatrix is a space of the constant negative curvature which value is equal to*

$$R_{\text{Ind}} = -\left(1 + \frac{1}{4}g^2\right) \le -1. \tag{A.29}$$

The respective *equation of* $\mathcal{E}_g^{SR}$*–geodesics* is of the form

$$\frac{d^2R^p}{ds^2} + C_q{}^p{}_r(g;R)\frac{dR^q}{ds}\frac{dR^r}{ds} = 0, \tag{A.30}$$

where s is the parameter of the arc-length defined in accordance with the rule

$$ds \overset{\text{def}}{=} \sqrt{|g_{pq}(g;R)dR^p dR^q|}. \tag{A.31}$$

The use of the functions

$$A(g;R) = R^0 - \frac{1}{2}g|\mathbf{R}|, \qquad \hat{A}(g;P) = P_0 + \frac{1}{2}g|\mathbf{P}| \tag{A.32}$$

and

$$L(g;R) = |\mathbf{R}| - \frac{1}{2}gR^0, \qquad \hat{L}(g;P) = |\mathbf{P}| + \frac{1}{2}gP_0 \tag{A.33}$$

is often convenient in various calculations.

In the limit $g \to 0$ the considered space degenerates to the ordinary pseudoeuclidean case:

$$B\big|_{g=0} = -[(R^0)^2 - \mathbf{R}^2], \qquad \hat{B}\big|_{g=0} = -[(P_0)^2 - \mathbf{P}^2],$$

$$j\big|_{g=0} = \hat{j}\big|_{g=0} = 1, \qquad F\big|_{g=0} = \sqrt{|(R^0)^2 - \mathbf{R}^2|},$$

$$H\big|_{g=0} = \sqrt{|(P_0)^2 - \mathbf{P}^2|}, \qquad g_{pq}\big|_{g=0} = e_{pq},$$

$$g^{pq}\big|_{g=0} = e^{pq}, \qquad C_{pqr}\big|_{g=0} = 0, \qquad R_{\mathrm{Ind}}\big|_{g=0} = -1.$$

Since at $g = 0$ the space $\mathcal{E}_g^{SR}$ is pseudoeuclidean, then $\mathcal{I}_{g=0}$ is the ordinary unit hyperboloid.

The $\mathcal{E}_g^{RS}$*–space invariant orthonormal frames:*

$$e_q^{\ P} \ := \sigma_q^{\ r} f_r^{\ P}, \qquad e_P^{\ q} \ := \mu_r^{\ q} m_P^{\ r} \tag{A.34}$$

are constructed on the basis of the formulas (B.27)–(B.31). The reciprocity $e_q^{\ P} e_R^{\ q} = \delta_R^P$ holds. Calculations show that

$$e_q^{\ P} = \frac{1}{h}\sigma_q^{\ P} + \frac{h-1}{h}\sigma^P l_q/F, \qquad e_P^{\ q} = h\mu_P^{\ q} + (1-h)\sigma_P l^q/F, \tag{A.35}$$

and

$$g_{pq}(g;R) = e_p^{\ 0}(g;R) e_q^{\ 0}(g;R) - \sum_{c=1}^{3} e_p^{\ c}(g;R) e_q^{\ c}(g;R). \tag{A.36}$$

The $\mathcal{E}_g^{RS}$*–space Ricci rotation coefficients*

$$R^{PQ}{}_p(g;t) \ := \left(\partial_i e_j^{\ Q} - C_p{}^r{}_s e_r^{\ Q}\right) e_T^{\ s} e^{TP} \tag{A.37}$$

can be obtained as the transform

$$R^{PQ}{}_p(g;R) = R^{PQ}{}_i(g;t)\sigma_p^{\ i} \tag{A.38}$$

from the quasi-pseudoeuclidean space. After using the formula (B.33) and (B.34), we find explicitly the components

$$R^{PQ}{}_q(g;R) = (h-1)(\sigma^P e_q^{\ Q} - \sigma^Q e_q^{\ P})/F^2(g;R), \tag{A.39}$$

or in another form

$$R^{PQ}{}_q(g;R) = \frac{h-1}{h}(\sigma^P \sigma_q^{\ Q} - \sigma^Q \sigma_q^{\ P})/F^2(g;R). \tag{A.40}$$

We have

$$\sigma^P/K = L^P, \qquad e_P^{\ q} L^P = l^q, \qquad e_P^{\ q} \sigma^P = R^q, \qquad R^{PQ}{}_p R^p = 0, \tag{A.41}$$

and

$$e_{Pt} e_{Qs}(\partial_p R^{PQ}{}_q - \partial_q R^{PQ}{}_p) = \frac{2(h-1)h}{F^2}(h_{pt}h_{qs} - h_{ps}h_{qt}) = \frac{2h}{1+h} S_{pqts}, \tag{A.42}$$

where $e_{Pt} = g_{rt}e^r_P$. Also,

$$\partial_p R^{PQ}{}_q - \partial_q R^{PQ}{}_p =$$

$$\frac{2(h-1)h}{F^2}\Big[e^P_p e^Q_q - e^Q_p e^P_q + \frac{1}{F}(-e^P_p\sigma^Q l_q - e^Q_q\sigma^P l_p + e^Q_p\sigma^P l_q + e^P_q\sigma^Q l_p)\Big] \tag{A.43}$$

and

$$R_{rsp} = (h-1)(l_r h_{sp} - l_s h_{rp})/F. \tag{A.44}$$

It will be noted that the frames (A.34) don't obey the "kinematic"property $e^p_0 \parallel R^p$. Instead, we have

$$e^P_q|_{g=0} = \delta^P_q, \qquad e^q_P|_{g=0} = \delta^q_P. \tag{A.45}$$

Appendix B. Quasi–pseudoeuclidean transformation

Let us introduce the nonlinear transformation

$$t^i = \sigma^i(g;R) \tag{B.1}$$

with the functions

$$\sigma^0 = \left|\frac{R^0 + g_-|\mathbf{R}|}{R^0 + g_+|\mathbf{R}|}\right|^{-G/4}\left(R^0 - \frac{1}{2}g|\mathbf{R}|\right), \qquad \sigma^a = h\left|\frac{R^0 + g_-|\mathbf{R}|}{R^0 + g_+|\mathbf{R}|}\right|^{-G/4} R^a; \tag{B.2}$$

$i, j, ... = 0, 1, 2, 3$ and $a, b, ... = 1, 2, 3$. With the help of the transformation, we can go over from the variables $\{R^p\}$ to the new variables $\{t^i\}$. The inverse transformation

$$R^p = \mu^p(g;t) \tag{B.3}$$

involves the functions

$$\mu^0 = \left|\frac{t^0 - m}{t^0 + m}\right|^{G/4}(t^0 + \frac{1}{2}Gm), \qquad \mu^a = \frac{1}{h}\left|\frac{t^0 - m}{t^0 + m}\right|^{G/4} t^a, \tag{B.4}$$

so that

$$t^i \equiv \sigma^i\left(g; \mu(g;t)\right), \qquad R^p \equiv \mu^p\left(g; \sigma(g;R)\right). \tag{B.5}$$

The notation

$$m = \sqrt{|e_{ab}t^a t^b|} \in [0, \infty)$$

has been used; the constant h is given by the formula (A.4).

Let us introduce the pseudoeuclidean metric function

$$S(t) \overset{\text{def}}{=} \sqrt{|e_{ij}t^i t^j|} \equiv \sqrt{|(t^0)^2 - m^2|} \tag{B.6}$$

($e_{ij} = \mathrm{diag}(1, -1, -1, -1)$ — the pseudoeuclidean metric tensor; $e^{ij} = e_{ij}$). It can readily be verified that the insertion of the functions (B.2) in (B.6) yields the identity

$$F(g;R) = S(t) \tag{B.7}$$

with the function $F(g;R)$ which is exactly the FMF (A.12). In this way, we call (B.1)–(B.2) the *quasi-pseudoeuclidean transformation.*

The functions (B.2) are obviously homogeneous of degree 1 with respect to the variable R:

$$\sigma^i(g;bR) = b\sigma^i(g;R), \qquad b>0, \tag{B.8}$$

from which it ensues that the derivatives

$$t^i_p(g;R) \overset{\text{def}}{=} \frac{\partial\sigma^i(g;R)}{\partial R^p} \equiv \sigma^i_p(g;R) \tag{B.9}$$

obey the identity

$$t^i_p(g;R)R^p = t^i. \tag{B.10}$$

Calculating the determinant gives merely

$$\det(t^i_p) = j^4h^3. \tag{B.11}$$

Similarly,

$$\mu^p(g;bt) = b\mu^p(g;t), \qquad b>0, \tag{B.12}$$

$$\mu^p_i(g;t) \overset{\text{def}}{=} \frac{\partial\mu^p(g;t)}{\partial t^i} \equiv R^p_i, \tag{B.13}$$

and

$$\mu^p_i(g;t)t^i = R^p. \tag{B.14}$$

Next, let us now construct the tensor

$$n^{ij}(g;t) \overset{\text{def}}{=} t^i_p t^j_q g^{pq}. \tag{B.15}$$

Straightforward rather lengthy calculations result in the following simple representations

$$n^{ij}(g;t) = h^2e^{ij} - \frac{1}{4}g^2l^il^j, \qquad n_{ij}(g;t) = \frac{1}{h^2}e_{ij} + \frac{1}{4}G^2l_il_j \tag{B.16}$$

($e_{ij}e^{jm} = \delta^m_i$ and $n_{ij}n^{jm} = \delta^m_i$), where

$$l^i \overset{\text{def}}{=} t^i/S(t), \qquad l_i \overset{\text{def}}{=} e_{ij}l^j \tag{B.17}$$

are respective pseudoeuclidean unit vectors. For them the equalities

$$l^il_i = 1, \quad n_{ij}l^j = l_i, \quad n^{ij}l_j = l^i, \quad n_{ij}l^il^j = 1, \quad n_{ij}t^it^j = S^2$$

are valid. The inversion of (B.15) can be written in the form

$$g_{pq}(g;R) = n_{ij}\left(g;\sigma(g;R)\right)t^i_p(g;R)t^j_q(g;R). \tag{B.18}$$

We also obtain

$$\det(n_{ij}) = -h^6. \tag{B.19}$$

We call the tensor $\{n_{ij}\}$ with the components (B.16) *quasi-pseudoeuclidean metric tensor*, and the very space

$$\mathcal{K}_g := \{V_4;\ S(t);\ n_{ij}(g;t);\ t\in V_4\} \tag{B.20}$$

quasi-pseudoeuclidean space. The formulas (B.7) and (B.15) show explicitly that the space defined is quasi-pseudoeuclidean image of the pseudo-Finsleroid relativistic space $\mathcal{E}_g^{SR}$, such that *when using the quasi-pseudoeuclidean transformations the studied pseudo-Finsleroid relativistic space* $\mathcal{E}_g^{SR}$ *transforms in the quasi-pseudoeuclidean space* $\mathcal{K}_g = \sigma(\mathcal{F}_g)$ *differed essentially from the pseudoeuclidean space* $\mathcal{E} \equiv \mathcal{K}_{g=0}$.

Let us evaluate from the tensor (B.16) the associated Christoffel symbols

$$N_m{}^i{}_n = n^{mk} N_{ikj}, \qquad N_{ikj} = \frac{1}{2}(n_{ik,j} + n_{jk,i} - n_{ij,k}). \tag{B.21}$$

We have subsequently

$$n_{ik,j} \overset{\text{def}}{=} \frac{\partial n_{ik}}{\partial t^j} = \frac{1}{4} g^2 (H_{ij} L_k + H_{kj} L_i)/S, \tag{B.22}$$

$$H_{mi} = e_{mi} - L_m L_i \equiv h^2 (n_{mi} - L_m L_i), \tag{B.23}$$

$$L^i H_{ij} = 0, \tag{B.24}$$

$$N_{mjn} = \frac{1}{4} G^2 H_{mn} L_j/S, \qquad N_m{}^i{}_n = \frac{1}{4} G^2 H_{mn} L^i/S, \tag{B.25}$$

and

$$N_i{}^m{}_j(t) = \frac{1}{4} G^2 L^m H_{ij}/S. \tag{B.26}$$

This entails the properties

$$t^i N_i{}^m{}_j = 0, \qquad N_i{}^j{}_j = 0.$$

The *quasi-pseudoeuclidean orthonormal frames* $\{m_P^i\}$ and $\{f_i^P\}$ subjected to the conditions

$$n^{ij} = e^{PQ} m_P^i m_Q^j, \qquad n_{ij} = e_{PQ} f_i^P f_j^Q, \tag{B.27}$$

prove to be taken in the simple form

$$f_i^P(g;t) = \frac{1}{h} h_j^P + \frac{h-1}{h} L_j L^P \tag{B.28}$$

and

$$m_P^j(g;t) = h h_P^j + (1-h) L_P L^j \tag{B.29}$$

with the vectors

$$L^P = h_j^P L^j, \qquad L_P = h_P^j L_j. \tag{B.30}$$

Here, $\{h_j^P\}$ and $\{h_P^j\}$ — orthonormal frames for the pseudoeuclidean metric tensor, so that

$$e^{ij} = e^{PQ} h_P^i h_Q^j, \qquad e_{ij} = e_{PQ} h_i^P h_j^Q. \tag{B.31}$$

The identity

$$f_i^P(g;t) t^i = t^P \equiv h_j^P t^j \tag{B.32}$$

holds.

The *associated quasi-pseudoeuclidean Ricci rotation coefficients*

$$R^{PQ}{}_i(g;t) := \left(\partial_i f_j^Q - N_i{}^n{}_j f_n^Q\right) m_T^j e^{TP} \tag{B.33}$$

are found merely as

$$R^{PQ}{}_i(g;t) = (h-1)(L^P f_i^Q - L^Q f_i^P)/S(t). \tag{B.34}$$

Also,

$$\frac{1}{8}\gamma^i R^{PQ}{}_i(\gamma_P\gamma_Q - \gamma_Q\gamma_P) = \frac{1}{4}\gamma^i R^{PQ}{}_i\gamma_P\gamma_Q = -\frac{3}{2}(h-1)\frac{\gamma_j t^j}{S^2}. \tag{B.35}$$

PROPOSITION.*The quasi-pseudoeuclidean metric tensor* $\{n_{ij}\}$ *is conformal to the pseudoeuclidean metric tensor. The conformal multiplier is equal to* ξ^{-2} *with*

$$\xi(g;t) = \left[\frac{1}{2}S^2(t)\right]^{\gamma/2}, \tag{B.36}$$

where

$$\gamma = h-1. \tag{B.37}$$

Indeed, applying the transformation

$$r^i = \xi(g;t)t^i/h \tag{B.38}$$

and using the coefficients

$$k^i_j \ := \frac{\partial r^i}{\partial t^j} = (\xi\delta^i_j + \xi' t^i t_j)/h = \xi\left(\delta^i_j + \gamma\frac{t^i t_j}{S^2(t)}\right)/h \tag{B.39}$$

to define the tensor

$$c^{ij}(g;r) \ := k^i_n k^j_m n^{nm}(g;t), \tag{B.40}$$

we obtain merely

$$c^{ij} = \xi^2 e^{ij}. \tag{B.41}$$

Therefore, the tensor

$$s_{pq}(g;R) = [\varkappa(g;R)]^2 g_{pq}(g;R) \tag{B.42}$$

meaningful over the initial $\mathcal{E}^{SR}_g$–space is conformally flat, where

$$\varkappa(g;R) = \left[\frac{1}{2}F^2(g;R)\right]^{\gamma/2} \equiv \xi(g;t). \tag{B.43}$$

The functions

$$\rho^i(g;R) = \frac{1}{h}\sigma^i(g;R)\varkappa(g;R) \equiv r^i \tag{B.44}$$

realize the transition from the initial FMT $\{g_{pq}\}$ to the pseudoeuclidean tensor $\{e_{ij}\}$. The inverse functions will be denoted as

$$\eta^p = \eta^p(g;\rho) \tag{B.45}$$

so that

$$\eta^p(g;\rho(g;R)) = R^p. \tag{B.46}$$

This entails the identities

$$\eta^p_m\rho^m_q=\delta^p_q,\qquad \eta^p_m\rho^n_p=\delta^n_m, \tag{B.47}$$

for the derivatives

$$\rho^m_p=\frac{\partial\rho^m}{\partial R^p} \tag{B.48}$$

and

$$\eta^p_m=\frac{\partial\eta^p}{\partial\rho^m}. \tag{B.49}$$

$$s_{pq}(g;R)=\rho^m_p(g;R)\rho^k_q(g;R)e_{mk} \tag{B.50}$$

The $\mathcal{E}^{SR}_g$–space *wave phase*

$$\Phi\ :=\frac{1}{h}k_nt^n\xi \tag{B.51}$$

obeys the important identity

$$\frac{\partial\Phi}{\partial t^j}=\xi f^P_j k_p \tag{B.52}$$

which entails directly that

$$n^{ij}\frac{\partial\Phi}{\partial t^i}\frac{\partial\Phi}{\partial t^j}=\xi^2e^{ij}k_ik_j \tag{B.53}$$

(see (B.27)). Inserting the quasi-pseudoeuclidean transformations (B.2) in (B.35) yields the representation

$$\Phi=k_n\rho^n(g;R), \tag{B.54}$$

so that

$$\Phi(g;R^0,R^1,R^2,R^3)=j\varkappa(\frac{1}{h}Ak_0-k_aR^a). \tag{B.55}$$

At the $O_1(g)$–level of consideration the phase (B.55) reduces to read

$$\Phi\approx\left(1-\frac{1}{4}g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|}\right)k_pR^p-\frac{1}{2}g|\mathbf{R}|k_0, \tag{B.56}$$

where $S^2=(R^0)^2-|\mathbf{R}|^2$.

Appendix C. Zero-phase conformally flat fronts in $\mathcal{E}^{SR}_g$–space

The zero-phase is described by the equation

$$\Phi(g;R^0,R^1,R^2,R^3)=0 \tag{C.1}$$

which, in view of (B.55), can explicitly be written as

$$(R^0-\frac{1}{2}g|\mathbf{R}|)k_0-hk_aR^a=0. \tag{C.2}$$

The last equation determines the *zero-phase wave surface,* which is obviously non-plane unless $g=0$. If for definiteness we specify the components $\{k_0,k_1,k_2,k_3\}$ to have $k_2=k_3=0$ in accordance with (5.13), the equation (C.2) reduces to merely

$$R^0-\frac{1}{2}g|\mathbf{R}|-hR^1=0, \tag{C.3}$$

or explicitly,

$$R^0 = hR^1 + \frac{1}{2}g|\mathbf{R}|. \tag{C.4}$$

Therefore, the point of intersection of the wave with the R^1–axis

$$R^1 > 0, \quad R^2 = R^3 = 0 \tag{C.5}$$

is governing by the equation

$$R^1 = g_+ R^0 \tag{C.6}$$

(see the definition of g_+ in the list (A.5)).

Thus we have

PROPOSITION. *The velocity of the point of intersection of the zero-phase surface with the R^1–axis is isotropic in accordance with* (C.6). *The vertex moves along the axis with the constant velocity*

$$v = g_+. \tag{C.7}$$

It will be noted that under the particular conditions (C.5) the vanishing $F = 0$ for the $\mathcal{E}_g^{SR}$–space implies the precise isotropic value (C.6) (see (A.12)).

Let us put

$$R_\perp = \sqrt{(R^2)^2 + (R^3)^2} \tag{C.8}$$

obtaining

$$|\mathbf{R}| = \sqrt{(R^1)^2 + (R_\perp)^2}. \tag{C.9}$$

We find

$$R^0 = \text{const} : \qquad \frac{\partial R^1}{\partial R_\perp} = -\frac{g}{2}\frac{R_\perp}{h|\mathbf{R}| + \frac{g}{2}R^1}, \tag{C.10}$$

so that

$$R^0 = \text{const} : \qquad \left.\frac{\partial R^1}{\partial R_\perp}\right|_{R_\perp = 0} = 0 \tag{C.11}$$

and

$$R^0 = \text{const} : \qquad \left.\frac{\partial^2 R^1}{\partial R_\perp^2}\right|_{R_\perp = 0} = -\frac{g}{2}g_+\frac{1}{R^1}. \tag{C.12}$$

The equation (11.4) can be resolved for R^1:

$$R^1 = hR^0 - \frac{g}{2}\sqrt{(R^0)^2 + (R_\perp)^2}. \tag{C.13}$$

The *asymptotic cone* $\mathcal{C}$ is determined by the equation

$$R^1 = -\frac{g}{2}R_\perp \tag{C.14}$$

and simultaneously presents the front at $R^0 = 0$. The asymptotic cone $\mathcal{C}$ is the same for any case of zero-phase wave. When $R^0 \to \infty$, the right-hand part in (C.12) tends to zero and, therefore, *the front is flattening around the vertex.*

In the approximation $O_1(g)$, we put $h = 1$ and reduce the formulae (C.2) and (C.4) to merely

$$(R^0 - \frac{1}{2}g|\mathbf{R}|)k_0 - k_a R^a = 0 \tag{C.15}$$

and

$$R^0 = R^1 + \frac{1}{2} g |\mathbf{R}|. \tag{C.16}$$

The equality (C.6) takes on the form

$$R^1 = (1 - \frac{1}{2} g) R^0, \tag{C.17}$$

and instead of (C.13) we get

$$R^1 = R^0 - \frac{g}{2} \sqrt{(R^0)^2 + (R_\perp)^2}. \tag{C.18}$$

Below, the zero-phase wave fronts determined by the equation (C.1)–(C.6) are shown at various circumstances.



Рис. 1: The fronts are pictured at the conventional pseudoeuclidean case, which is the trivial



Рис. 2: The case $g = 2$



Рис. 3: The fronts are shown for the value $g = 1.2$. The higher case pictures correspond



Рис. 4: The fronts are shown for the value $g = -1.2$. The higher case pictures correspond





Рис. 5: The case $g = 0.8$; $R^0 = 25$ at the top picture and $R^0 = -25$ at the bottom picture



Рис. 6: With $R^0 = -60, -45, -30, -15, 0, 15, 30, 45, 60,$ the fronts are simulated in the two-

We preluded with Fig. 1 which symbolized the ordinary pseudoeuclidean plane wave, that is, the $(g = 0)$–case.

In Fig. 2 the fronts were depicted at various values of the parameter $g \neq 0$, showing that the fronts are non-plane. In the higher figures the values $R^0 = 0; 15; 30; 45$ were subsequently be taken. Lower figures symbolized the front at the initial momentum $R^0 = 25$ and the arrow shows the direction of propagation. At $R^0 = 0$ the vertex is a conic point. We observe how the front is flattening in a vicinity of vertex as soon as the value of R^0 is increasing.

In Figs. 3 and 4 the fronts were shown separately at the values $R^0 = 0, 50, 100, 150$ for positive and definite values of g. Additionally, Fig. 5 shows two kindred fronts at two opposite values of the temporal coordinate R^0. Finally, Fig. 6 showed the succession of the fronts in the $2d$-case where the vanishing $R^3 = 0$ has been assumed.

When $g < 0$ the fronts move in the direction pointed by the vertex; the opposite direction of motion takes place at $g > 0$. In each figure the fronts move in the positive direction of the R^1–axis.

Appendix D. Conformal flatness of the space $\mathcal{E}_g^{SR}$

Let us use the scalar

$$\mathcal{K}(g; R) = \left(\frac{1}{2}F^2(g; R)\right)^{h-1} \equiv \bigl(\varkappa(g; R)\bigr)^2 \tag{D.1}$$

and introduce the new tensor $s_{pq}(g; R)$ according to

$$g_{pq}(g; R) = \frac{1}{\mathcal{K}(g; R)} s_{pq}(g; R). \tag{D.2}$$

Taking the associated Christoffel symbols

$$S_p{}^t{}_r = \frac{1}{2} s^{ts} \left(\frac{\partial s_{ps}}{\partial R^r} + \frac{\partial s_{rs}}{\partial R^p} - \frac{\partial s_{pr}}{\partial R^s} \right), \tag{D.3}$$

we can straightforwardly evaluate the curvature tensor

$$M_t{}^u{}_{rs} \overset{\text{def}}{=} \frac{\partial S_t{}^u{}_r}{\partial R^s} - \frac{\partial S_t{}^u{}_s}{\partial R^r} + S_t{}^w{}_r S_w{}^u{}_s - S_t{}^w{}_s S_w{}^u{}_r. \tag{D.4}$$

The result proves to be zero:

$$M_t{}^u{}_{rs} \equiv 0. \tag{D.5}$$

Thus we get

PROPOSITION. *The space $\mathcal{E}_g^{SR}$ is conformally flat. The conformal multiplier is a function of the FMF according to* (D.1)–(D.2).

Therefore, the transformation

$$r^i = \rho^i(g; R) \tag{D.6}$$

should exist such that in terms of the coefficients

$$\rho^i_q(g; R) \overset{\text{def}}{=} \frac{\partial \rho^i(g; R)}{\partial R^q} \tag{D.7}$$

the identity

$$\frac{1}{h^2}s_{pq}(g;R)=\rho^i_p(g;R)\rho^j_q(g;R)e_{ij} \tag{D.8}$$

holds fine ($e_{ij}=\mathrm{diag}(1,-1,-1,-1)$ is a pseudoeuclidean metric tensor). A careful consideration leads to

PROPOSITION. *The conformal transformation* (D.6) *is given by the components*

$$\rho^0(g;R)=\frac{1}{h}\varkappa(g;R)j(g;R)\left(R^0-\frac{1}{2}g|\mathbf{R}|\right),\quad \rho^a(g;R)=\varkappa(g;R)j(g;R)R^a. \tag{D.9}$$

The identity

$$\left(R^0-\frac{1}{2}g|\mathbf{R}|\right)^2-h^2|\mathbf{R}|^2=-B(g;R) \tag{D.10}$$

is useful to apply when verifying this Proposition.

The transformation inverted to (D.6) is given by the law

$$R^p=\eta^p(g;r) \tag{D.11}$$

with the functions

$$\eta^0(g;r)=\nu(g;r)\left(hr^0+\frac{1}{2}g|\mathbf{r}|\right)/j(g;r),\quad \eta^a(g;r)=\nu(g;r)r^a/j(g;r), \tag{D.12}$$

where

$$\nu(g;r)=\left(\frac{h}{\sqrt{2}}S(g;r)\right)^{(h-1)/h}=\frac{1}{\varkappa(g;R)} \tag{D.13}$$

and

$$j(g;r)=\left(\frac{r^0-|\mathbf{r}|}{r^0+|\mathbf{r}|}\right)^{-G/4}. \tag{D.14}$$

Using the derivatives

$$\eta^p_i(g;r)\overset{\text{def}}{=}\frac{\partial\eta^p(g;r)}{\partial r^i}, \tag{D.15}$$

we may rewrite (D.8) in the backward direction as follows:

$$\eta^p_i(g;r)\eta^q_j(g;r)s_{pq}(g;R)=e_{ij}. \tag{D.16}$$

It is convenient to use the notation

$$\gamma=h-1. \tag{D.17}$$

In the time-like case we get

$$\rho^0(g;R)=\frac{1}{h}\varkappa(g;R)j(g;w)E(g;w)R^0,\quad \rho^a(g;R)=\varkappa(g;R)j(g;w)R^a,\quad R\in S^+_g, \tag{D.18}$$

where

$$E(g;w)=1-\frac{1}{2}gw. \tag{D.19}$$

The function j entering (D.37) has been defined by (A.13). We obtain the identity

$$\Big(E(g;w)\Big)^2 - h^2w^2 = Q(g;w) \tag{D.20}$$

and the representations

$$\rho^0_0(g;R) = \Big(\gamma E(1-gw) + Q - \frac{1}{2}gwE\Big)\,\frac{j\varkappa}{Qh^2}, \tag{D.21}$$

$$\rho^0_a(g;R) = \Big(-\gamma E + \frac{g(E-Q)}{2w}\Big)\,\frac{j\varkappa w_a}{Qh^2}, \tag{D.22}$$

$$\rho^a_0(g;R) = \Big(\gamma(1-gw) - \frac{1}{2}gw\Big)\,\frac{j\varkappa w^a}{Qh}, \tag{D.23}$$

$$\rho^a_b(g;R) = \Big(-\gamma w_b w^a + Q\delta^a_b + \frac{gw_bw^a}{2w}\Big)\,\frac{j\varkappa}{Qh}, \tag{D.24}$$

from which the identity

$$\delta_{cd}\rho^c_a\rho^d_b = \left(\delta_{ab} + \frac{w_aw_b}{Qw}\Big(-\gamma w + \frac{1}{2}g\Big)\Big(2 - \frac{\gamma w^2}{Q} + \frac{gw}{2Q}\Big)\right)\frac{(j\varkappa)^2}{h^2} \tag{D.25}$$

follows. Some simplification is applicable to find

$$\rho^0_0(g;R) = \big(E(1-gw) - h\,w^2\big)\,\frac{j\varkappa}{Qh}, \tag{D.26}$$

$$\rho^0_a(g;R) = (h-E)\,\frac{j\varkappa w_a}{Qh}, \tag{D.27}$$

$$\rho^a_0(g;R) = \big(hQ - E + h\,w^2\big)\,\frac{j\varkappa w^a}{Qh}, \tag{D.28}$$

$$\rho^a_b(g;R) = \left(Q\delta^a_b + \Big((1-h)w + \frac{g}{2}\Big)\,\frac{w_bw^a}{w}\right)\frac{j\varkappa}{Qh}, \tag{D.29}$$

together with

$$\delta_{cd}\rho^c_a(g;R)\rho^d_b(g;R) = \left[\delta_{ab} + \Big(g + (1-h)w + (h^2-h)\frac{w(1+w^2)}{Q}\Big)\,\frac{w_aw_b}{Qw}\right](j\varkappa)^2. \tag{D.30}$$

Also,

$$\det(\rho^i_p) = (j\varkappa)^N, \qquad \det(\rho^a_b) = \big(E - h\,w^2\big)\,\frac{(j\varkappa)^{N-1}}{Q}, \tag{D.31}$$

$$w^b \rho_b^a = \left(E - h\, w^2\right) \frac{j \varkappa w^a}{Q}, \qquad w_a \rho_b^a = \left(E - h\, w^2\right) \frac{j \varkappa w_b}{Q}, \tag{D.32}$$

$$\partial_q \varkappa = \frac{\gamma \varkappa}{F^2} R_q \tag{D.33}$$

$(N = 4)$.

For the reciprocal coefficients defined in accordance with (D.15) we find the representations

$$\eta_0^0(g; r) = \left(E - h\, w^2\right) \frac{h}{j \varkappa Q}, \tag{D.34}$$

$$\eta_a^0(g; r) = (E - h)\, \frac{w_a h}{j \varkappa Q}, \tag{D.35}$$

$$\eta_0^a(g; r) = \left(E - h w^2 - h Q\right) \frac{w^a h}{j \varkappa Q}, \tag{D.36}$$

$$\eta_b^a(g; r) = \left(Q \delta_b^a + \left((E - h) w - \frac{1}{2} g Q \right) \frac{w_b w^a}{w} \right) \frac{h}{j \varkappa Q}, \tag{D.37}$$

and the identities

$$R_p \eta_0^p = \frac{E F^2 h}{R^0 j \varkappa Q}, \qquad R_p \eta_a^p = -\frac{h^2 F^2 w_a}{R^0 j \varkappa Q}, \tag{D.38}$$

$$\frac{\partial \left(\varkappa^{-3} \sqrt{|\det(s_{rs})|}\, \eta_0^p \right)}{\partial R^p} = -6 \gamma \frac{E F^2 j^3 h}{R^0 Q}. \tag{D.39}$$

In the space-like region,

$$\rho^0(g; R) = \frac{1}{h} \varkappa(g; R) j(g; k) f(g; k) q, \quad \rho^a(g; R) = \varkappa(g; R) j(g; k) R^a, \qquad R \in \mathcal{R}_g^+, \tag{D.40}$$

where

$$f(g; k) = k - \frac{1}{2} g, \tag{D.41}$$

so that

$$h^2 - \left(f(g; k) \right)^2 = L(g; k) \tag{D.42}$$

and

$$\rho_0^0(g; R) = \left(-\gamma (k - g) f + L + \frac{1}{2} g f \right) \frac{j \varkappa}{L h}, \tag{D.43}$$

$$\rho_a^0(g;R) = \Big(\gamma f - \frac{1}{2}g(L+kf)\Big)\frac{j\varkappa n_a}{Lh}, \tag{D.44}$$

$$\rho_0^a(g;R) = \Big(-\gamma(k-g) + \frac{1}{2}g\Big)\frac{j\varkappa n^a}{L}, \tag{D.45}$$

$$\rho_b^a(g;R) = \Big(L\delta_b^a + (\gamma - \frac{1}{2}gk)n^a n_b\Big)\frac{j\varkappa}{L} \tag{D.46}$$

or

$$\rho_0^0(g;R) = \Big(h - (k-g)f\Big)\frac{j\varkappa}{L}, \tag{D.47}$$

$$\rho_a^0(g;R) = \Big(f - kh\Big)\frac{j\varkappa n_a}{L}, \tag{D.48}$$

$$\rho_0^a(g;R) = \Big(f - (k-g)h\Big)\frac{j\varkappa n^a}{L}, \tag{D.49}$$

$$\rho_b^a(g;R) = \Big(L\delta_b^a + (h - 1 - \frac{1}{2}gk)n^a n_b\Big)\frac{j\varkappa}{L}, \tag{D.50}$$

$$\det(\rho_p^i) = (j\varkappa)^N, \qquad \det(\rho_b^a) = (h - fk)\,\frac{(j\varkappa)^{N-1}}{L} \tag{D.51}$$

$(N = 4)$.

Also,

$$n^b\rho_b^a = (h - fk)\,\frac{j\varkappa n^a}{L}, \qquad w_a\rho_b^a = (h - fk)\,\frac{j\varkappa n_b}{L}, \tag{D.52}$$

$$\partial_q\varkappa = -\frac{\gamma\varkappa}{F^2}R_q. \tag{D.53}$$

The reciprocal coefficients are obtainable to read

$$\eta_0^0(g;r) = (h - fk)\,\frac{1}{j\varkappa L}, \tag{D.54}$$

$$\eta_a^0(g;r) = (kh - f)\,\frac{n_a}{j\varkappa L}, \tag{D.55}$$

$$\eta_0^a(g;r) = ((k-g)h - f)\,\frac{n^a}{j\varkappa L}, \tag{D.56}$$

$$\eta^a_b(g;r) = \left(L\delta^a_b + (h-(k-g)f-L)\,n_b n^a\right)\frac{1}{j\varkappa L}, \tag{D.57}$$

entailing

$$R_p\eta^p_0 = -\frac{fF^2}{qj\varkappa L}, \qquad R_p\eta^p_a = \frac{hF^2 n_a}{qj\varkappa L}, \tag{D.58}$$

$$\frac{\partial\left(\varkappa^{-3}\sqrt{|\det(s_{rs})|}\,\eta^p_0\right)}{\partial R^p} = 6\gamma\frac{fF^2j^3}{qL}. \tag{D.59}$$

Appendix E. $O_1(g)$–approximations

Considering the parameter g to be small:

$$|g| \ll 1 \tag{E.1}$$

and performing expansions with respect to the parameter, we shall denote by the symbol $O_1(g)$ the terms proportional to the first degree of g and by $O_{\geq 2}(g)$ — the terms proportional to the second and higher degrees of g.

First of all, the respective approximation of the components of the FMT reads as follows:

$$g_{00}(g;R) = 1 - g\frac{|\mathbf{R}|R^0}{(R^0)^2-|\mathbf{R}|^2} - \frac{1}{2}g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|} + O_{\geq 2}(g), \tag{E.2}$$

$$g_{0a}(g;R) = g\frac{|\mathbf{R}|R^a}{(R^0)^2-|\mathbf{R}|^2} + O_{\geq 2}(g), \tag{E.3}$$

$$g_{ab}(g;R) = -\delta_{ab} + \frac{1}{2}g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|}\delta_{ab} - g\frac{R^aR^bR^0}{|\mathbf{R}|((R^0)^2-|\mathbf{R}|^2)} + O_{\geq 2}(g); \tag{E.4}$$

$$g^{00}(g;R) = 1 + g\frac{|\mathbf{R}|R^0}{(R^0)^2-|\mathbf{R}|^2} + \frac{1}{2}g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|} + O_{\geq 2}(g), \tag{E.5}$$

$$g^{0a}(g;R) = g\frac{|\mathbf{R}|R^a}{(R^0)^2-|\mathbf{R}|^2} + O_{\geq 2}(g), \tag{E.6}$$

$$g^{ab}(g;R) = -\delta^{ab} - \frac{1}{2}g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|}\delta^{ab} + g\frac{R^aR^bR^0}{|\mathbf{R}|((R^0)^2-|\mathbf{R}|^2)} + O_{\geq 2}(g). \tag{E.7}$$

The occurrence of the logarithmic function is characteristic of the approximations. For the function (A.15) we get

$$j(g;R) = 1 - \frac{1}{4}g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|} + O_{\geq 2}(g) \tag{E.8}$$

so that, owing to (A.18),

$$\det(g_{pq}) = -1 + 2g\ln\frac{R^0-|\mathbf{R}|}{R^0+|\mathbf{R}|} + O_{\geq 2}(g). \tag{E.9}$$

The structure of the tensor C_{pqr} is such that the tensor is proportional to g, so that

$$C_{pqr} = gS_{pqr} + O_{\geq 2}(g), \tag{E.10}$$

where the components S_{pqr} are given by the list

$$R^0 S_{000} = \frac{R^0|\mathbf{R}|^3}{((R^0)^2 - |\mathbf{R}|^2)^2}, \qquad R^0 S_{a00} = -\frac{(R^0)^2|\mathbf{R}|R^a}{((R^0)^2 - |\mathbf{R}|^2)^2}, \tag{E.11}$$

$$R^0 S_{ab0} = \frac{1}{2}\frac{R^0|\mathbf{R}|}{(R^0)^2 - |\mathbf{R}|^2}\delta_{ab} + \frac{1}{2}\frac{R^0R^aR^b((R^0)^2 + |\mathbf{R}|^2)}{|\mathbf{R}|((R^0)^2 - |\mathbf{R}|^2)^2}, \tag{E.12}$$

$$R^0 S_{abc} = -\frac{1}{2}\frac{(R^0)^2}{|\mathbf{R}|((R^0)^2 - |\mathbf{R}|^2)}(\delta_{ab}R^c + \delta_{ac}R^b + \delta_{bc}R^a) + \frac{(R^0)^2R^aR^bR^c((R^0)^2 - 3|\mathbf{R}|^2)}{2|\mathbf{R}|^3((R^0)^2 - |\mathbf{R}|^2)^2}. \tag{E.13}$$

For the components

$$S_p{}^q{}_r = e^{qs}S_{psr}, \qquad S^{pqr} = e^{ps}e^{qt}e^{rv}S_{ptv}, \tag{E.14}$$

we obviously obtain

$$S_0{}^0{}_0 = S_{000}, \qquad S_0{}^a{}_0 = -S_{0a0}, \qquad S_a{}^0{}_0 = -S_{a00}, \qquad S_a{}^b{}_c = -S_{abc}, \tag{E.15}$$

and

$$S^{000} = S_{000}, \qquad S^{a00} = -S_{a00}, \qquad S^{ab0} = S_{ab0}, \qquad S^{abc} = -S_{abc}. \tag{E.16}$$

For contractions $C_p = C_p{}^q{}_q$ we get

$$C_p = gS_p + O_{\geq 2}(g), \quad C^p = gS^p + O_{\geq 2}(g), \tag{E.17}$$

and

$$R^0 S_0 = -\frac{N}{2}\frac{R^0|\mathbf{R}|}{(R^0)^2 - |\mathbf{R}|^2}, \qquad R^0 S_a = \frac{N}{2}\frac{(R^0)^2R^a}{|\mathbf{R}|((R^0)^2 - |\mathbf{R}|^2)}, \tag{E.18}$$

$$S^0 = S_0, \qquad S^a = -S_a. \tag{E.19}$$

Obviously

$$C_pC^p = O_{\geq 2}(g). \tag{E.20}$$

Also,

$$\partial_p C^p = -\frac{N}{2}g(N-2)\frac{R^0}{|\mathbf{R}|((R^0)^2 - |\mathbf{R}|^2)} \tag{E.21}$$

$((N = 4))$.

The above corrections are singular over the light cone, that is, when $R^0 - |\mathbf{R}| = 0$. The field $\{S_p\}$ introduces a preferred vector over the cone. The tensor $\{S_{pqr}\}$ assigns an anisotropic structure over the background pseudoeuclidean space and enters respective covariant derivatives.

At the $O_1(g)$–level of consideration the curvature tensor of the space under study is zero:

$$S_{pqrs} = 0 + O_{\geq 2}(g) \tag{E.22}$$

(see (A.22) and (A.23)). The same conclusion is applicable to the quasi-pseudoeuclidean objects:

$$n_{ij}(g;t) = e_{ij} + O_{\geq 2}(g), \qquad n^{ij} = e^{ij} + O_{\geq 2}(g), \tag{E.23}$$

$$N_i{}^k{}_j(g;t) = 0 + O_{\geq 2}(g), \qquad R_{mnij}(g;t) = 0 + O_{\geq 2}(g) \tag{E.24}$$

(see (B.16), (B.26), and (B.34)). Also, the conformal properties are trivial at the $O_1(g)$–level of consideration:

$$\gamma = O_{\geq 2}(g) \tag{E.25}$$

(see (B.37)).

References

[1] H. Busemann: *Canad. J. Math.* **1** (1949), 279.

[2] H. Rund: *The Differential Geometry of Finsler spaces,* Springer-Verlag, Berlin 1959.

[3] G.S. Asanov: *Rep. Math. Phys.* **45** (2000), 155; **47** (2001), 323.

[4] G.S. Asanov:arXiv:hep-ph/0306023, 2003; arXiv:math.MG/0402013, 2004; arXiv:math-ph/0406029, 2004.

[5] G.S. Asanov: *Publ. Math. Debrecen* **61** (2005), 1.

[6] G.S. Asanov:arXiv:gr-qc/0204070, 2002.